

# New dialysis studies inform delivery of care, ways to improve patient outcomes

November 10 2022

---



Credit: Pixabay/CC0 Public Domain

In the U.S., dialysis is a costly treatment with poor health outcomes. According to the Centers for Disease Control and Prevention, 15% of adults—37 million people—have chronic kidney disease. As the disease

progresses to end-stage kidney disease, patients are required to receive dialysis regularly or have a kidney transplant. Although dialysis is lifesaving in the short-term, five-year mortality rates exceed 60%.

Eugene Lin, senior fellow at the USC Schaeffer Center for Health Policy & Economics and assistant professor of nephrology at the Keck School of Medicine of USC, is conducting research to improve the way we think about [dialysis care](#) and the outcomes for patients.

"Dialysis is unique because it is the only disease-qualifying condition for Medicare. Because it is a multibillion-dollar industry that impacts a disproportionate portion of Medicare spending, understanding the policies that affect the provision of dialysis and downstream outcomes is critical," said Lin.

In studies published in November 2022, Lin and his colleagues analyze two distinct aspects of dialysis and their effect on patient outcomes: Ownership of dialysis facilities and timing of dialysis and surgery.

## **Dialysis facilities owned by kidney doctors may provide better care**

Clinics that are entirely or partially owned by nephrologists represent an increasing share of dialysis facilities. Unlike nephrologists that do not have an ownership stake in a dialysis clinic, physician owners can directly profit from the types of treatments they recommend. But they may also have more autonomy in terms of clinical decisions.

In a study published in *JAMA Internal Medicine*, Lin and his colleagues analyzed data from fee-for-service Medicare patients receiving dialysis for end-stage kidney disease from January to November in 2017. The researchers compared outcomes between patients managed by physician

owners and patients managed by non-owners within clinics owned by physicians.

Patients treated by physician-owners were approximately 25% more likely to receive home dialysis care. They were also less likely to use erythropoietin-stimulating agents (ESAs), which are expensive medications that are used to treat anemia in patients with end-stage kidney disease. Strikingly, less use of these drugs was not associated with worse patient outcomes with no difference in blood transfusions.

On measures of missed appointments, transplant wait lists, deaths, hospitalizations or long-term catheter use, the researchers find physician owners performed no better or worse than non-physician owner clinics.

"It may be that dialysis clinics that are owned by nephrologists develop different clinical guidelines—they may be better equipped to build successful home dialysis programs or develop more sophisticated triggers for anemia so that they do not have to prescribe these drugs as often," explained Lin.

"We had thought that the financial conflict of interest stemming from clinic ownership would lead to potentially worse outcomes for patients," Lin said. "This doesn't seem to happen. However, our study was limited to a cross-sectional analysis of 2017 data. We need better transparency of financial relationships across the entire dialysis industry, so that policymakers can conduct more robust studies of these conflicts of interest."

## **A shorter interval between dialysis and surgery may improve post-surgery survival**

The optimal timing for having dialysis prior to elective surgery is

unknown. But these patients have an increased risk of post-operative mortality, [heart attack](#), stroke sepsis and surgical site infection and the timing of when they receive dialysis may play an important role.

In the second study, published in *JAMA*, Lin and colleagues analyze data on 90-day post operative mortality for end-stage kidney patients 18 years of age and older who underwent elective surgery from 2011 to 2018. Of the 1.1 million surgical procedures included in the data, 65% of procedures had a 1-day interval between dialysis and surgery while 36% of procedures had either a 2- or 3-day interval. For 16.6% of procedures, dialysis was performed on the day of the surgery.

The researchers find longer intervals between dialysis and surgery were associated with a higher risk of post-operative mortality. For example, among patients with a 3-day interval, the mortality risk was 5.2% compared to 4.2% for patients with 1-day interval. Yet, having dialysis treatment on the day of the surgery erased the increased risk.

"Given our findings, doctors and hospitals should consider scheduling surgeries around dialysis or, when that is not possible, perform dialysis right before the surgery," said Lin, who was the senior author on this study.

The researchers note that while the risk difference was small—and many of the procedures included in the study were relatively minor—patients with end-stage kidney disease undergo a lot of procedures and generally face worse outcomes, so even small changes in the absolute risk could be meaningful.

## **Policy implications**

Given the resources required for dialysis and the poor health outcomes experienced by patients with end-stage kidney disease, Lin suggests that

policymakers and stakeholders consider a number of policy changes.

- Improved data transparency would allow researchers to better understand whether financial incentives and ownership are impacting patient outcomes. Regardless of physician-ownership status, several recent [studies](#) conducted by researchers at the Schaeffer Center have identified aspects of the dialysis market that are leading to increased spending.
- CMS could establish quality metrics on post-operative outcomes for patients with end-stage kidney disease undergoing surgery or on the scheduling of elective surgeries.

"Ultimately, I hope these projects facilitate improvements in dialysis delivery, so that patients on [dialysis](#) receive the best possible care," Lin said.

**More information:** Eugene Lin et al, Association Between Nephrologist Ownership of Dialysis Facilities and Clinical Outcomes, *JAMA Internal Medicine* (2022). [DOI: 10.1001/jamainternmed.2022.5002](https://doi.org/10.1001/jamainternmed.2022.5002)

Vikram Fielding-Singh et al, Association Between Preoperative Hemodialysis Timing and Postoperative Mortality in Patients With End-stage Kidney Disease, *JAMA* (2022). [DOI: 10.1001/jama.2022.19626](https://doi.org/10.1001/jama.2022.19626)

Provided by Keck School of Medicine of USC

Citation: New dialysis studies inform delivery of care, ways to improve patient outcomes (2022, November 10) retrieved 8 May 2024 from <https://medicalxpress.com/news/2022-11-dialysis-delivery-ways-patient-outcomes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.