

Evidence supports three-year interval for multi-target stool DNA screening for those at average risk of colon cancer

November 29 2022



Credit: Unsplash/CC0 Public Domain

A scientific study exploring the appropriate interval for colorectal cancer screening via non-invasive multi-target stool DNA testing for individuals

with average risk for the disease reported finding no colorectal cancers three years after an initial negative multi-target stool DNA test. These results suggest that at least a three-year interval between screenings using this method is clinically appropriate.

The results of this multicenter study, led by Regenstrief Institute Research Scientist and Indiana University School of Medicine faculty member Thomas F. Imperiale, M.D., were determined via repeating the multi-target stool DNA test at year three, with test results confirmed via colonoscopy, the gold standard test for detecting colon cancers.

While no cancers were found by screening at the three-year interval, 63 advanced precancerous lesions were identified in the 591 study participants. The study authors note that because the test targets cancer and high risk pre-cancerous polyps and is not designed to identify non-advanced lesions, some found at three years may have become advanced since the previous multi-target stool DNA screening.

"This study provides screening age range adults and their clinicians with evidence-based information that they need on multi-target stool DNA screening frequency," said Dr. Imperiale.

"However, due to the constraints imposed by the pandemic when individuals and healthcare systems postponed or canceled appointments, the number of participants dwindled over the three years, and further study of the test interval is required. A three-year interval may be the most appropriate, but it is possible that a longer interval may work as well or better."

A study of 10,000 patients led by Dr. Imperiale, published in the *New England Journal of Medicine* in 2014, reported that multi-target stool DNA testing, which requires a single stool sample expelled from the body directly into a container, detects 92.3 percent of colon cancers.

The ability of a test to detect disease—known as sensitivity—is the most important characteristic for cancer screening tests because the primary role of these tests is to "rule out" disease—in this case, a cancer of the colon or rectum. In 2021, national guidelines lowered the age recommended to begin screening for colon cancer from 50 to 45.

"Colon cancer screening tests like the multi-target stool DNA or the less sensitive annual fecal immunochemical test [FIT] are efficient ways to screen the population, especially those at the low-risk end of the average risk population, which includes most individuals in the younger age range of screening," said Dr. Imperiale.

"Using at-home tests encourages those who appreciate their ease of use and lower hassle factor to be screened. And it doesn't simply benefit the individual, it benefits others who are high risk and are best screened with colonoscopy."

Approximately seven or eight out of 10 individuals who fall within the range of those for whom [colorectal cancer screening](#) is recommended by national guidelines are considered to be at average risk for the disease.

Colon cancer is the third most common cancer in the U.S. and has a significant death rate for both men and women.

The findings are published in the journal *Cancer Prevention Research*.

More information: Thomas F. Imperiale et al, Three-Year Interval for the Multi-Target Stool DNA Test for Colorectal Cancer Screening: A Longitudinal Study, *Cancer Prevention Research* (2022). [DOI: 10.1158/1940-6207.CAPR-22-0238](#)

Thomas F. Imperiale et al, Multitarget Stool DNA Testing for Colorectal-Cancer Screening, *New England Journal of Medicine* (2014). [DOI:](#)

[10.1056/NEJMoa1311194](https://doi.org/10.1056/NEJMoa1311194)

Provided by Regenstrief Institute

Citation: Evidence supports three-year interval for multi-target stool DNA screening for those at average risk of colon cancer (2022, November 29) retrieved 3 May 2024 from

<https://medicalxpress.com/news/2022-11-evidence-three-year-interval-multi-target-stool.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.