


H7N9 avian influenza in humans: Sex differences in disease outcomes

November 16 2022



Correlation of testosterone and cytokine/chemokine levels in men infected with H7N9 influenza. a–g Shown is the observed regression between testosterone and cytokine/chemokine levels in H7N9 IAV-infected males. Natural logarithm transformed (Ln) testosterone expression levels measured in the sera from H7N9 males were plotted over the Ln transformed cytokine/chemokines levels. a–e Overall regression of testosterone with cytokines IL-6 (interleukin 6; a), IL-10 (interleukin 10; b), IL-15 (interleukin 15; c), IL-1RA (interleukin 1 receptor antagonist; d) and IL-1 α (interleukin 1 alpha; e) observed from H7N9 younger males without interaction between deaths and survivors (18–49 yrs survival/death: n = 18/15); f, g Observed overall regression of testosterone with cytokines IL-6 (interleukin 6; f) and IL-10 (interleukin 10; g) in H7N9 older males without interaction between deaths and survivors (≥ 50 yrs survival/death: n = 27/11). a–g The best-fit line with 95% confidence intervals of the overall regression is shown in each figure. The center of the error distribution is the regression line. Two-tailed linear regression was performed. Adjustment for multiple hypotheses was not necessary due to the explorative nature of the study. A hollow blue circle represents survival, and a solid blue circle represents death. R squared values are shown in the figures. Credit: *Nature Communications* (2022). DOI: 10.1038/s41467-022-34500-5

An interdisciplinary team from the Leibniz Institute of Virology and China CDC shows for the first time that infection with the H7N9 avian influenza virus attacks the hormonal axis of males but not females.

In their study published in *Nature Communications*, the scientists retrospectively analyzed one of the largest H7N9 influenza cohorts and showed that low testosterone levels are associated with the development of severe or even fatal disease in males. In contrast, seasonal H1N1 or H3N2 influenza did not significantly affect the hormonal axis in male or [female patients](#). This study identifies low testosterone levels caused by avian H7N9 infection as a poor prognostic marker in males.

Prior to the COVID-19 pandemic, H7N9 avian influenza viruses were

considered to be the viruses with the highest risk of causing the next pandemic. In March 2013, [avian influenza virus](#) type A (H7N9) crossed species boundaries for the first time in Eastern China, jumping from birds to humans and showing [higher incidence](#) in men compared to women. In the following five epidemic waves, the H7N9 incidence was repeatedly higher in men than women.

To unravel the mechanisms behind these [gender differences](#), the German-Chinese team analyzed patients with laboratory-confirmed H7N9 infection and compared to H7N9-negative close contacts as well as patients with seasonal influenza infection. The scientists show that H7N9 infection specifically attacks the hormone axis in men but not women.

In men, H7N9 infection mediates [low testosterone levels](#) that correlates with severe and even fatal outcomes. In murine models, the authors further confirm the [causal link](#) between H7N9 infection and testosterone depletion in males. They further show that avian H7N9 influenza virus replicates in the testes of mice and induces local and systemic inflammation that likely interferes with testosterone production.

Blueprint for other respiratory virus infections

To date, very little is known about the molecular mechanisms that lead to sex-specific disease courses in respiratory virus infections. This study could serve as a blueprint for investigating sex differences in other respiratory infections, including SARS-CoV-2, as observed in the current pandemic.

"Avian influenza viruses continue to pose a high epidemic and pandemic risk. The 2021/2022 season was the largest avian influenza epidemic recorded worldwide, including Europe. Thus, understanding [molecular mechanisms](#) that mediate sex-specific disease outcomes are crucial for individualized patient management," explains Prof. Gülşah Gabriel, head

of LIV's Viral Zoonoses–One Health research unit and professor at the University of Veterinary Medicine Hannover.

"The strict surveillance and massive vaccination in poultry have so far prevented H7N9 virus further spread to humans. But avian influenza viruses continue to evolve and require high vigilance," says Prof. Yuelong Shu, the former director of Chinese National Influenza Center at China CDC.

More information: Tian Bai et al, H7N9 avian influenza virus infection in men is associated with testosterone depletion, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-34500-5](https://doi.org/10.1038/s41467-022-34500-5)

Provided by Leibniz-Institut für Virologie (LIV)

Citation: H7N9 avian influenza in humans: Sex differences in disease outcomes (2022, November 16) retrieved 23 May 2024 from <https://medicalxpress.com/news/2022-11-h7n9-avian-influenza-humans-sex.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--