


# MRI reveals significant brain abnormalities post-COVID

November 21 2022



Group analysis on susceptibility-weighted imaging exhibiting higher susceptibility-weighted imaging values in the COVID group when compared to healthy controls. Three significant clusters were found primarily in the white matter regions of the pre-frontal cortex and in the brainstem. The clusters (a) and (b) are observed bilaterally in the cerebral white matter near the orbitofrontal gyrus, whereas (c) lies in the midbrain region. Credit: RSNA and Sapna S. Mishra

Using a special type of MRI, researchers have uncovered brain changes in patients up to six months after they recovered from COVID-19, according to a study being presented next week at the [annual meeting](#) of the Radiological Society of North America (RSNA).

About one in five adults will develop [long-term effects](#) from COVID-19, according to the U.S. Centers for Disease Control and Prevention. Neurological symptoms associated with long COVID include difficulty thinking or concentrating, headache, sleep problems, lightheadedness, pins-and-needles sensation, change in smell or taste, and depression or anxiety. However, studies have found that COVID-19 may be associated with changes to the heart, lungs or other organs even in asymptomatic patients.

As more people become infected and recover from COVID-19, research has begun to emerge, focusing on the lasting consequences of the disease.

For this study, researchers used susceptibility-weighted imaging to analyze the effects that COVID-19 has on the brain. Magnetic susceptibility denotes how much certain materials, such as blood, iron and calcium, will become magnetized in an applied magnetic field. This ability aids in the detection and monitoring of a host of neurologic conditions including microbleeds, vascular malformations, brain tumors and stroke.

"Group-level studies have not previously focused on COVID-19 changes in [magnetic susceptibility](#) of the brain despite several case reports signaling such abnormalities," said study co-author Sapna S. Mishra, a Ph.D. candidate at the Indian Institute of Technology in Delhi. "Our study highlights this new aspect of the neurological effects of COVID-19 and reports significant abnormalities in COVID survivors."

The researchers analyzed the susceptibility-weighted imaging data of 46 COVID-recovered patients and 30 healthy controls. Imaging was done within six months of recovery. Among patients with long COVID, the most commonly reported symptoms were fatigue, trouble sleeping, lack of attention and memory issues.

"Changes in susceptibility values of brain regions may be indicative of local compositional changes," Mishra said. "Susceptibilities may reflect the presence of abnormal quantities of paramagnetic compounds, whereas lower susceptibility could be caused by abnormalities like calcification or lack of paramagnetic molecules containing iron."

MRI results showed that patients who recovered from COVID-19 had significantly higher susceptibility values in the frontal lobe and brain stem compared to healthy controls. The clusters obtained in the frontal lobe primarily show differences in the white matter.

"These [brain regions](#) are linked with fatigue, insomnia, anxiety, depression, headaches and cognitive problems," Mishra said.

Portions of the left orbital-inferior frontal gyrus (a key region for language comprehension and production) and right orbital-inferior frontal gyrus (associated with various cognitive functions including attention, motor inhibition and imagery, as well as social cognitive processes) and the adjacent [white matter](#) areas made up the [frontal lobe](#) clusters.

The researchers also found a significant difference in the right ventral diencephalon region of the brain stem. This region is associated with many crucial bodily functions, including coordinating with the endocrine system to release hormones, relaying sensory and motor signals to the [cerebral cortex](#) and regulating [circadian rhythms](#) (the sleep-wake cycle).

"This study points to serious long-term complications that may be caused by the coronavirus, even months after recovery from the infection," Mishra said. "The present findings are from the small temporal window. However, the longitudinal time points across a couple of years will elucidate if there exists any permanent change."

The researchers are conducting a longitudinal study on the same patient cohort to determine whether these brain abnormalities persist over a longer time frame.

Co-authors are Rakibul Hafiz, Ph.D., Tapan Gandhi, Ph.D., Vidur Mahajan, M.B.B.S., Alok Prasad, M.D., and Bharat Biswal, Ph.D.

Provided by Radiological Society of North America

Citation: MRI reveals significant brain abnormalities post-COVID (2022, November 21)  
retrieved 8 May 2024 from  
<https://medicalxpress.com/news/2022-11-mri-reveals-significant-brain-abnormalities.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--