

Neuronal mechanism involved in the learning of maternal behavior discovered

November 8 2022



Credit: AI-generated image ([disclaimer](#))

Various conditions such as postpartum depression or postpartum psychosis can lead to an alteration in maternal behavior and disrupt the mother-child bonding process. A research team has conducted a study in which they were able to identify the neuronal circuits in the brain that are activated during the learning of maternal behavior.

The findings, published in *The EMBO Journal*, can serve as a basis for developing therapeutic interventions.

In the course of their preclinical investigation, the scientists led by Daniela Pollak from the Division of Neurophysiology and Neuropharmacology at MedUni Vienna's Center for Physiology and Pharmacology analyzed maternal behavior of female mice towards newborn pups. While the neural processes involved in the development of maternal care behavior in female mice after birth have already been described, the current study addressed the question of which circuits in the brain are activated during the learning of care behavior in nulliparous virgin females, [female mice](#) that have not been pregnant.

The researchers found the answer in the [anterior cingulate cortex](#) (ACC), a region in the prefrontal lobe of the brain that is associated, among other things, with the recognition and evaluation of social processes and the development of emotional awareness. The [mouse model](#) showed that in females who have not been pregnant or given birth, the ACC gets activated when they acquire maternal behavior upon first contact with pups.

"Our observations have demonstrated that, through repeated experience with pups, the virgin females are capable of learning maternal behavior that fully resembles those of mothers after delivery," explains project leader Pollak, reporting from her research. The study revealed that during this learning process ACC activity is controlled by an excitatory feedback circuit involving a specific group of neurons in a central brain region (the thalamus).

Learning through repeated experience

Maternal behavior is defined, among other things, by sensitivity and responsiveness to signals of infant needs. It is instinctively displayed in

virtually all mammals and upon first contact with newborns immediately after delivery. In some species, such as rodents, even animals that have never given birth may exhibit maternal care behavior toward newborns. In this case, the care behavior, such as returning displaced pups from outside the nest back into the nest area, where they are warm and protected from predators, are acquired through repeated experience with the pups.

We know, for example from observations in [adoptive parents](#), that humans can also learn parental behaviors. Various pathological conditions, such as [postpartum depression](#) or [postpartum psychosis](#) can lead to alterations in maternal behavior and disrupt the mother-child bonding process.

"By showing that maternal behaviors can be acquired and identifying the underlying [neuronal circuits](#) in the [brain](#) that control this acquisition, we are creating a potential basis for developing therapeutic options for these clinical situations," says Pollak, outlining the translational relevance of the study results, which were obtained in collaboration with Tibor Harkany from the Division of Molecular Neurosciences at MedUni Vienna's Center for Brain Research.

More information: Micaela Glat et al, An accessory prefrontal cortex–thalamus circuit sculpts maternal behavior in virgin female mice, *The EMBO Journal* (2022). [DOI: 10.15252/emboj.2022111648](https://doi.org/10.15252/emboj.2022111648)

Provided by Medical University of Vienna

Citation: Neuronal mechanism involved in the learning of maternal behavior discovered (2022, November 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-11-neuronal-mechanism-involved-maternal-behavior.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.