

Personalized kidney screening for people with type 1 diabetes could reduce costs, detect disease earlier

November 2 2022



Credit: Pixabay/CC0 Public Domain

Taking a personalized approach to kidney disease screening for people with type 1 diabetes (T1D) may reduce the time that chronic kidney

disease (CKD) goes undetected, according to a new analysis performed by the Epidemiology of Diabetes Interventions and Complications study group.

The finding, published in *Diabetes Care*, provides the basis for the first evidence-based kidney screening model for people with T1D.

Current CKD screening recommendations include annual urinary albumin excretion rate (AER) testing for anyone who has had T1D for at least five years. Albumin is a protein found in the blood and having too much albumin in the urine is a sign of kidney disease.


The new findings suggest that AER screening could be personalized to optimize testing frequency and early detection of CKD. Specifically, people with T1D who are at low risk of developing CKD could be tested for AER less frequently to reduce burden and cost, and those at high risk for CKD could be tested more frequently to facilitate earlier CKD detection.

People with T1D have an estimated 50% risk of developing CKD over their lifetime. CKD can progress to [kidney failure](#), requiring dialysis or a kidney transplant. Using more than 30 years of participant data of AER and HbA1c (an integrated measure of blood glucose) from 1,334 participants in the NIDDK-funded Diabetes Control and Complications Trial (DCCT) and the observational follow-up Epidemiology of Diabetes Interventions and Complications (EDIC) study, the [study group](#) identified three levels of CKD risk that were associated with a later CKD diagnosis. They then developed a model to estimate the optimal screening intervals for people with T1D to detect CKD at its earliest stages.

Optimal Frequency of Urinary Albumin Screening in Type 1 Diabetes
 Bruce A. Perkins, Ionut Bebu, Ian H. de Boer, Mark Molitch, Bernard Zinman,
 John Bantle, Gayle M. Lorenzi, David M. Nathan, John M. Lachin

A PERSONALIZED ALTERNATIVE TO ANNUAL SCREENING IN TYPE 1 DIABETES CAN SUBSTANTIALLY
 REDUCE BOTH THE TIME WITH UNDETECTED KIDNEY DISEASE AND THE FREQUENCY OF
 URINE TESTING.

<ul style="list-style-type: none"> • Kidney disease screening recommendations include annual urine testing for albuminuria after 5 years' duration of type 1 diabetes. • We aimed to determine a simple, risk factor-based screening schedule that optimizes early detection and testing frequency. 	<ul style="list-style-type: none"> • Urinary albumin excretion measurements were obtained from 1,343 participants in the DCCT/EDIC study. • Likelihood of the onset of moderately or severely elevated albuminuria (confirmed albumin excretion rate [AER] ≥ 30 mg/24 h) and its risk factors were used to identify individualized screening schedules. 	<p>Screening in 2 years for those with current AER ≤ 10 mg/24 h and HbA1c $\leq 8\%$ (0.6% 3-year cumulative incidence), in 6 months for those with AER 21-30 mg/24 h or HbA1c $\geq 9\%$ (8.9% 3-year cumulative incidence), and 1 year for all others (2.4% 3-year cumulative incidence) was associated with 34.9% reduction in time with undetected albuminuria and 20.4% reduction testing frequency as compared with annual screening.</p>
---	---	---

Diabetes Care Funding: Division of Diabetes, Endocrinology & Metabolic Diseases of the National Institute of Diabetes and Digestive and Kidney Disease (current grant numbers U01 DK094176 and U01 DK094157)  American Diabetes Association. Connected for Life.

Credit: *Diabetes Care* (2022). DOI: 10.2337/dc22-1420

According to the model's findings:

- People with AER of 21–30 mg per 24 hours and a HbA1c of at least 9% are at high risk for developing CKD and could be screened for urine albumin every six months. This screening frequency could reduce time with undetected kidney disease so that appropriate interventions can be instituted as early as possible.
- Those with AER ≤ 10 mg per 24 hours and a HbA1c $\leq 8\%$ are at lower risk for developing CKD and could be screened every two years. This change reduces patient burden and potentially saves millions of dollars compared to annual [screening](#).
- All others with T1D \geq five years could continue to be screened annually.

The DCCT, which took place from 1983 to 1993, found that, for people

with T1D, keeping [blood glucose levels](#) close to normal greatly reduced the chances of developing eye, [kidney](#), and nerve disease. Its follow-up study, EDIC, began in 1994 to explore how [diabetes](#) affects the body over time and the long-term benefits of early and intensive blood glucose control in the development of later diabetes complications.

"Optimal Frequency of Urinary Albumin Screening in Type 1 Diabetes" appears Nov. 2 in *Diabetes Care*.

More information: Bruce A. Perkins et al, Optimal Frequency of Urinary Albumin Screening in Type 1 Diabetes, *Diabetes Care* (2022). [DOI: 10.2337/dc22-1420](https://doi.org/10.2337/dc22-1420)

Provided by National Institutes of Health

Citation: Personalized kidney screening for people with type 1 diabetes could reduce costs, detect disease earlier (2022, November 2) retrieved 23 June 2024 from <https://medicalxpress.com/news/2022-11-personalized-kidney-screening-people-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.