


# Study finds risk of seizures is higher after COVID-19 than after influenza

November 16 2022

---



Credit: Pixabay/CC0 Public Domain

People who have a COVID-19 infection are more likely to develop seizures or epilepsy within the next six months than people who have an influenza infection, according to a study published in the November 16,

2022, online issue of *Neurology*. The increased risk was more noticeable in children than adults. It was also more noticeable in people who did not need hospitalization for COVID-19 infections.

"While the overall risk of developing [seizures](#) or [epilepsy](#) was low—less than 1% of all people with COVID-19, given the large number of people who have been infected with COVID-19, this could result in increases in the number of people with seizures and epilepsy," said study author Arjune Sen, MD, Ph.D., of the University of Oxford in England. "In addition, the increased risk of seizures and epilepsy in children gives us another reason to try to prevent COVID-19 infections in kids."

For the study, researchers looked at a [health records](#) network for people with COVID-19 infections. They were matched to people who were diagnosed with influenza during the same time period and who were similar in age, sex and other factors, such as other [medical conditions](#). None of the participants had previously been diagnosed with epilepsy or recurrent seizures. The researchers then looked to see whether people developed epilepsy or seizures in the following six months.

There were 152,754 people in each of the COVID-19 and influenza groups.

People who had COVID-19 were 55% more likely to develop epilepsy or seizures over the next six months than people who had influenza. The rate of new cases of epilepsy or seizures was 0.94% in the people who had COVID-19, compared to 0.60% in those who had influenza.

"People should interpret these results cautiously since the overall risk is low," Sen said. "We do, however, recommend that [health care professionals](#) pay particular attention to individuals who may have more subtle features of seizures, such as focal aware seizures, where people are alert and aware of what is going on, especially in the three months

following a less severe COVID-19 infection."

A limitation of the study was that researchers were unable to identify which specific virus variants people were infected with, which could have influenced results.

**More information:** Maxime Taquet et al, Incidence of Epilepsy and Seizures Over the First 6 Months After a COVID-19 Diagnosis: A Retrospective Cohort Study, *Neurology* (2022). DOI: 10.1212/WNL.000000000000201595 , [dx.doi.org/10.1212/WNL.000000000000201595](https://doi.org/10.1212/WNL.000000000000201595)

Provided by American Academy of Neurology

Citation: Study finds risk of seizures is higher after COVID-19 than after influenza (2022, November 16) retrieved 2 May 2024 from <https://medicalxpress.com/news/2022-11-seizures-higher-covid-influenza.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------