

AI improves colorectal cancer screening in Lynch syndrome

December 28 2022, by Inka Våth



(from left) Prof. Jacob Nattermann and Dr. Robert Hüneburg in the endoscopy.
Credit: Rolf Müller, University Hospital Bonn (UKB)

People with Lynch syndrome have a higher hereditary risk of colon cancer. Despite regular endoscopic surveillance, it remains elevated in

those affected. Researchers at the National Center for Hereditary Tumor Diseases (NZET) at Bonn University Hospital (UKB) have now found that artificial intelligence (AI) can improve the effectiveness of colonoscopy in the presence of Lynch syndrome. The study results have been published online in the *United European Gastroenterology Journal*.

Hereditary non-polyposis colorectal cancer (HNPCC)—Lynch syndrome (LS) for short—is the most common hereditary colorectal cancer syndrome. Based on estimates, it is believed to affect approximately 300,000 individuals in Germany alone. LS is responsible for about 2–3% of all colon cancers. It is triggered by defects in genes responsible for repairing the human genetic material DNA. Children of those affected have a 50% risk of also having the pathologically altered hereditary disposition, technically known as a mutation, and thus also a high risk of developing colon cancer at a young age.

Therefore, regular monitoring by means of [colonoscopy](#) is recommended here every one to two years. "However, despite such regular endoscopic monitoring, the risk of colorectal cancer remains elevated in those affected," says Prof. Jacob Nattermann, head of the Hepatogastroenterology Section at UKB's Medical Clinic I. The researchers explain this by the fact that colonoscopy still misses a considerable number of adenomas, the potential precursors of colorectal [cancer](#). "Small and flat adenomas in particular are at risk of being overlooked even by experienced gastroenterologists," says Prof. Nattermann.

Increased adenoma detection rate

Recent data suggest that [artificial intelligence](#) (AI)-assisted colonoscopy, also known as computer-aided detection (CADe), may help increase adenoma detection rates (ADR) in the general population.

"Therefore, the aim of our study was to evaluate the diagnostic performance of AI-assisted colonoscopy in patients with Lynch [syndrome](#)," says Robert Hüneburg, senior physician at UKB's Medical Clinic I.

In close collaboration with the Institute of Medical Informatics, Statistics and Epidemiology at the University of Leipzig, 46 LS patients were studied using standard endoscopy and 50 LS patients with AI-assisted colonoscopy between December 2021 and December 2022. Here, significantly more of the AI-assisted examinations (36%) detected adenomas than the standard examinations (26.1%).

"This was mainly due to significantly improved detection of flat adenomas," Hüneburg says.

This study was the first in the world to collect data suggesting that AI-assisted real-time colonoscopy is a promising approach to optimize endoscopic surveillance of LS patients, especially to improve detection of flat [adenomas](#). "Due to the [small sample size](#), based on these results, a larger multicenter study is now planned under the leadership of our department," said Prof. Nattermann.

More information: Robert Hüneburg et al, Real-time use of artificial intelligence (CADEYE) in colorectal cancer surveillance of patients with Lynch syndrome—A randomized controlled pilot trial (CADLY), *United European Gastroenterology Journal* (2022). [DOI: 10.1002/ueg2.12354](https://doi.org/10.1002/ueg2.12354)

Provided by University of Bonn

Citation: AI improves colorectal cancer screening in Lynch syndrome (2022, December 28) retrieved 17 April 2024 from

<https://medicalxpress.com/news/2022-12-ai-colorectal-cancer-screening-lynch.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.