


New clue discovered as to how and why cancer cells spread

December 5 2022, by Gillian Rutherford



Viscosity enhances cell migration and promotes an ARP2/3-mediated dense actin network at the leading edge. **a,b**, Speeds of MDA-MB-231 cells (**a**) and other indicated cell types (**b**) inside confining channels at prescribed viscosities. The red lines represent the median of ≥ 69 cells from ≥ 3 experiments. **c**, Cell trajectories on 2D collagen-coated surfaces after 10 h. **d**, Cells disseminating from 3D spheroids. **e**, The time required for the first cell dissociation from each

spheroid ($n \geq 53$) from 3 experiments. **f**, Airyscan images of phalloidin stained cells on collagen-coated substrates. The red arrow indicates high F-actin staining along the cell edge. **g**, The fraction of cell-projected area with a Lifeact–GFP-rich lamella for $n \geq 28$ cells from 3 experiments. **h**, The leading edge of Lifeact–GFP-expressing cells on collagen-coated surfaces at $t = 0$ min (red) and $t = 2$ min (cyan) (left). Right, leading-edge lamella growth in $n \geq 19$ cells from 3 experiments. Data are the moving average \pm s.e.m. *P*

Citation: New clue discovered as to how and why cancer cells spread (2022, December 5) retrieved 6 May 2024 from <https://medicalxpress.com/news/2022-12-clue-cancer-cells.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.