

# Cluster headaches may be more severe in women

December 21 2022

---



Credit: CC0 Public Domain

While cluster headache is more common in men than in women, a new study suggests that the disorder may be more severe for women. The study is published in the December 21, 2022, online issue of *Neurology*, the medical journal of the American Academy of Neurology.

Cluster headaches are short but extremely painful headaches that can occur many days, or even weeks, in a row. The headaches can last anywhere from 15 minutes to three hours.

"Cluster headache is still often misdiagnosed in women, perhaps because some aspects can be similar to migraine," said study author Andrea C. Belin, Ph.D., of the Karolinska Institutet in Stockholm, Sweden. "It is important for physicians to be aware of how the disorder manifests differently in men and women so the most effective treatment can be given as fast as possible."

The study involved 874 people diagnosed with [cluster](#) headaches, with 66% male and 34% female. Participants answered a detailed questionnaire about their symptoms, medications, headache triggers and lifestyle habits.

Women were more likely to be diagnosed with chronic cluster headache than men. Chronic cluster headache is defined as recurring cluster headache attacks for one year or more without interruption, or with short intermissions with no symptoms that last less than three months. Eighteen percent of women were diagnosed with chronic cluster headache, compared to 9% of men.

Attacks also lasted longer for women than for men. For example, 8% of women said headache bouts lasted an average of four to seven months, compared to 5% of men, while 26% of women said bouts on average lasted less than one month, compared to 30% of men.

Women were also more likely to report that their attacks occurred at various times throughout the day than men, 74% to 63%. Women were more likely to have a family member with a history of cluster headache, 15% to 7%.

"While the ratio of men to women with cluster headache has been shifting over the years, it is still considered mainly a disorder of men, making it more difficult for women with milder symptoms to be diagnosed with cluster [headache](#) than men," Belin said. "It's possible this could contribute to the higher rate of chronic [cluster headache](#) in [women](#)."

A limitation of the study was that information was reported by the participants, so they may not have remembered everything correctly.

**More information:** Mark J. Burish et al, Cluster Headache: Worse in Women, *Neurology* (2022). DOI: 10.1212/WNL.0000000000206807 , [dx.doi.org/10.1212/WNL.0000000000206807](https://doi.org/10.1212/WNL.0000000000206807)

Provided by American Academy of Neurology

Citation: Cluster headaches may be more severe in women (2022, December 21) retrieved 26 April 2024 from <https://medicalxpress.com/news/2022-12-cluster-headaches-severe-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.