


# COVID-19 virus can affect vision and depth perception, finds study

December 21 2022



Clinical features and virus titres in the eyes of SARS-CoV-2-infected mice. Eight- to nine-week-old male K18-hACE2 mice were intranasally mock-infected or infected with 10<sup>4</sup> PFU of SARS-CoV-2 (n = 5 for mock- and SARS-CoV-2-infected mice, respectively; Mock, Gray; SARS-CoV-2, Red). a Body weight changes shown as percentage of starting weight. b Survival was evaluated at the indicated dpi. c Representative image of tearing and eye discharge in SARS-CoV-2-infected mice at 6 dpi (right) compared to those in mock-infected mice (left). d Viral load in the lungs and eyes, including appendages, was

analyzed using a plaque assay at 6 dpi. e Viral RNA levels in the lungs, brain, eyes, including appendages, lacrimal gland, parotid gland and spleen were assessed using RT-qPCR at 6 dpi. Viral RNA copies were cut-off ( $10^4$  copies/ $\mu\text{g}$ ). A dashed line indicates the viral RNA levels of the spleen as a limit of detection. f Representative confocal images of immunofluorescence stained retinal sections of IN-infected mice ( $n = 3$  per group) for viral spike (S) protein (red) at 6 dpi. DAPI staining (blue) was used to visualize nuclei of the ganglion cell layer (GCL), inner nuclear layer (INL), and outer-nuclear layer (ONL) in the retinal cross-section. Scale bar = 100  $\mu\text{m}$ . Symbols represent means  $\pm$  SEM. Statistically significant differences between the groups were determined using multiple two-tailed t-tests (a), unpaired two-tailed t-test (d) or one-way ANOVA (e). SARS-CoV-2: severe acute respiratory syndrome coronavirus 2; PFU plaque-forming unit, vRNA viral RNA, dpi days post-infection. Credit: *Nature Communications* (2022). DOI: 10.1038/s41467-022-35225-1

Researchers are investigating whether the COVID virus can affect vision and depth-perception of those infected.

The study, co-led by Griffith University's Menzies Health Institute and South Korea's Center for Convergent Research for Emerging Virus Infection, Korea Research Institute of Chemical Technology, aims to understand how SARS-CoV-2 affects the eyes and whether it could serve as a virus infection route.

It found the eyes and the trigeminal nerves are susceptible to the virus and that (in animal models) SARS-CoV-2 can infect the eye through the [respiratory tract](#), via the brain.

Principal Research Leader and co-lead author Prof Suresh Mahalingam said the virus can begin to affect vision when inflammation of the optic nerves, abnormal fluid build-up, and immune cell infiltration cause the retina to get thicker.

"The virus can infect the eye through nerve tissues at the back of the eye that play a role in the visual aspects of the eye and sending signals for visual purposes," Prof Mahalingam said.

"The result of this retinal inflammation was a reduction in depth perception due to [blurred vision](#)."

This blurred [vision](#) does appear to be symptomatic only, not a permanent degeneration of the eye tissue.

It is also only likely to affect a very small number of people.

Griffith University Ph.D. student Mr. Ng Wern Hann said that while a lot of COVID research has been focused on respiratory infection, particularly in the lungs and nasal region, there has not been much focus on the eyes.

"We found the virus can indeed infect the eye through a normal intranasal approach, but also if droplets of the virus make direct contact with the eye," he said.

"The ACE2 receptor is what the [virus](#) attaches to in order to infect a particular cell in a tissue or organ, and this receptor is found in abundance in the lungs, tonsils, [nasal cavity](#), kidneys and heart, which is why a lot of reports have been published for those organs, but we found ACE2 receptors are also present in the eye, therefore facilitating [infection](#)."

The findings published in *Nature Communications* give new insights into COVID-19 disease and may facilitate the development of treatment strategies for patients.

**More information:** Gi Uk Jeong et al, Ocular tropism of SARS-

CoV-2 in animal models with retinal inflammation via neuronal invasion following intranasal inoculation, *Nature Communications* (2022). [DOI: 10.1038/s41467-022-35225-1](https://doi.org/10.1038/s41467-022-35225-1)

Provided by Griffith University

Citation: COVID-19 virus can affect vision and depth perception, finds study (2022, December 21) retrieved 25 April 2024 from <https://medicalxpress.com/news/2022-12-covid-virus-affect-vision-depth.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.