

December 16 2022, by Ingrid Fadelli

## Study finds that insomnia could cause lower back pain and vice versa

<figure>

Individual estimates about the potential causal effect of insomnia and low back pain. The x-axis shows that the single nucleotide polymorphism (SNP) effect on insomnia in (A) and the y-axis shows the SNP effect on low back pain; (B) shows the results of the reverse causal inference. Several methods of MR analysis including MR Egger, weighted median, inverse variance weighted (IVW), maximum likelihood(ML) and penalized weighted median were performed. Credit: *Frontiers in Neuroscience* (2022). DOI: 10.3389/fnins.2022.1074605



Throughout the course of their life, most humans are likely to experience some form of lower back pain. This pain is sometimes associated with injuries, chronic back issues, medical conditions, the menstrual cycle or pregnancy, and other underlying physiological causes. In many cases, however, pinpointing the origin of lower back pain is difficult, despite its severity and its potential impact on the day-to-day life of those affected.

Some studies have found associations between <u>lower back pain</u> and different psychological and <u>social factors</u>, including gender, body weight, smoking habits and mood disorders. More recently, research also linked this sometimes crippling experience to insomnia and <u>sleep</u> <u>disturbances</u>.

Researchers at Zhejiang University School of Medicine have recently carried out a study exploring these recent findings further, by trying to better understand the relationship between sleep and lower back pain. Their findings, published in *Frontiers in Neuroscience*, suggest that there is a bidirectional causal relationship between insomnia and lower back pain, with each potentially causing the other.

"Previous observational studies have shown that <u>low back pain</u> often coexists with sleep disturbances, however, the causal relationship remains unclear," Ge Luo, Yuanyuan Yao, Jiachun Tao, Tingting Wang and Min Yan wrote in their paper. "In the present study, the causal relationship between sleep disturbances and LBP was investigated and the importance of sleep improvement in the comprehensive management of LBP was emphasized."

Luo and his colleagues examined the self-reported and <u>genetic data</u> of over 400,000 people of European ancestry, collected as part of the



genome-wide association study (GWAS) by the UK Biobank. These people had completed genetic tests and answered a series of questions about their sleep patterns.

The researchers' analyses focused on different factors associated with poor sleep quality, including insomnia, long sleep duration, short sleep duration, and daytime sleepiness. They thus screened the GWAS data and selected participants with genetic variants associated with different dimensions of poor sleep.

To infer a possible causal relationship between poor sleep and lower back pain, they used Mendelian randomization (MR), a relatively new technique that allows scientists to use measured genetic variations to unveil possible genetic and environmental risk factors that could cause poor health or specific conditions.

Their findings highlighted a bidirectional <u>causal relationship</u> between insomnia and lower back pain, meaning that insomnia can cause lower back pain, and vice versa. In addition, they found that lower back pain could cause daytime sleepiness, although not the opposite.

In the future, these results could inspire further studies exploring the relationship between poor <u>sleep</u> and insomnia, potentially incorporating people with other ethnic and genetic backgrounds. In addition, the work by Luo and his colleagues could promote the exploration of therapeutic interventions for lower back pain that also target insomnia and unhealthy sleeping patterns.

"The main results of our research showed a potential bidirectional causal association of genetically predicted insomnia with lower back pain," Luo and his colleagues wrote in their paper. "Sleep improvement may thus be important in comprehensive management of LBP."



**More information:** Ge Luo et al, Causal association of sleep disturbances and low back pain: A bidirectional two-sample Mendelian randomization study, *Frontiers in Neuroscience* (2022). <u>DOI:</u> <u>10.3389/fnins.2022.1074605</u>

© 2022 Science X Network

Citation: Study finds that insomnia could cause lower back pain and vice versa (2022, December 16) retrieved 25 April 2024 from <u>https://medicalxpress.com/news/2022-12-insomnia-pain-vice-versa.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.