


New findings on neuronal activities in the sensorimotor cortex

December 5 2022



Neurons from layer 5 of the motor cortex stained with a fluorescent dye. Credit: Ilka Diester



An interdisciplinary research team at the University of Freiburg has found important clues about the functioning of the sensorimotor cortex. The new findings on neuronal activities in this brain area could be helpful for the further development and use of so-called neuroprostheses. These have an interface with the nervous system and are intended to help compensate for neuronal dysfunctions.

"Our results will contribute to the improvement of neuroprosthetic approaches while shortening the training period of patients with prostheses," says neurobiologist Prof. Dr. Ilka Diester from the Faculty of Biology at the University of Freiburg. The results have just been published in the journal *Nature Communications*.

Understanding the brain under more natural conditions

The research project also involved the working groups of computer scientist Prof. Dr. Thomas Brox from the University of Freiburg and neuroscientist Prof. Dr. Daniel Durstewitz from the Central Institute of Mental Health in Mannheim. The team found evidence of conserved structures of neuronal activity in the sensorimotor cortex of freely moving rats. The <u>electrophysiological recordings</u> across the entire bilateral sensorimotor cortex allow conclusions about the respective contributions of the premotor, motor and sensory areas. In particular, the researchers found a clear gradient for a contralateral bias, i.e., for movements of the opposite half of the body, from anterior to posterior regions.

Previous findings on sensorimotor cortex are mostly based on highly constrained, stereotyped movements in a laboratory setting. The current work uses recordings on freely moving subjects using 3D tracking and addresses the question of the transferability of knowledge about neural



control of movements from constrained behavior to a freely moving condition, a prerequisite for understanding the brain under more <u>natural</u> <u>conditions</u> as well as for the further development of neuroprosthetic devices.

Categories of behavior across individuals

The team used a dimensionality reduction and neural data alignment method. Thus, the high-dimensional neuronal patterns were reduced to a low-dimensional representation by means of their similarity to other patterns, resulting in geometric structures in the <u>visual representation</u>. These <u>geometric patterns</u> were then automatically aligned with each other, similar to the image of holding a magnet to a pack of nails. These then align themselves in a certain direction.

Based on such aligned <u>geometric structures</u>, the researchers were able to decipher behavioral categories across recording sessions and even across individuals, and find corresponding evidence of conserved structures of neural activity.

More information: Svenja Melbaum et al, Conserved structures of neural activity in sensorimotor cortex of freely moving rats allow cross-subject decoding, *Nature Communications* (2022). <u>DOI:</u> <u>10.1038/s41467-022-35115-6</u>

Provided by University of Freiburg

Citation: New findings on neuronal activities in the sensorimotor cortex (2022, December 5) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2022-12-neuronal-sensorimotor-cortex.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.