

Seizures seem tied to faster decline in people with dementia

December 2 2022, by Cara Murez



Dementia patients who suffer from seizures tend to decline faster and



die younger, according to a new study that urges caregivers to watch for these sudden brain changes.

"Our hope is that controlling <u>seizures</u> by prescribing antiseizure medications to these patients will slow down the progression of cognitive impairment," said <u>Dr. Ifrah Zawar</u>, lead study author and an assistant professor at the University of Virginia School of Medicine.

"Unfortunately, seizures are often underdiagnosed because they can be subtle and the person just seems confused, so family members often mistake them for typical signs of <u>dementia</u>," Zawar added in a news release from the American Epilepsy Society.

In some people, a staring spell is evidence of a seizure, while others may experience uncontrolled jerking movement of their arms or legs.

Researchers analyzed data from 2005 to 2021 from 39 Alzheimer's disease research centers, determining that 374 (1.4%) of the more than 26,000 dementia patients had seizures.

Those patients were significantly younger when mental decline began, under 63 years old compared to 68 years old. They also died younger, at just under 73 years old compared to nearly 80 years old.

The dementia patients who had seizures were more likely to have a genetic mutation related to Alzheimer's disease, to have suffered a stroke or <u>traumatic brain injury</u>, to have depression, or to have less education, the researchers found.

Dementia patients with seizures had more advanced cognitive impairment, including problems with thinking, communication, understanding and memory. They also had more difficulty with basic daily activities, including eating, bathing, dressing and using the



bathroom.

Investigators also found they were more likely to be dependent on others for physical help.

"We know dementia patients are more likely to have seizures, and that people who have seizures are more likely to develop dementia, but it's still a 'chicken and egg question' regarding which one causes the other," Zawar said.

"It's important for <u>family members</u> and <u>health care providers</u> to recognize the patients with dementia who are at high risk for developing seizures, and ensure they are diagnosed and treated in a timely fashion," she advised.

Seizures occur in as many as 64% of people who have dementia, the researchers said in background notes, and those with dementia are six to 10 times more likely to develop seizures.

The findings were scheduled for presentation Friday at the American Epilepsy Society <u>annual meeting</u>, in Nashville. Findings presented at medical meetings are considered preliminary until published in a peer-reviewed journal.

More information: The U.S. Centers for Disease Control and Prevention has more on <u>dementia</u>.

Copyright © 2022 HealthDay. All rights reserved.

Citation: Seizures seem tied to faster decline in people with dementia (2022, December 2) retrieved 10 May 2024 from https://medicalxpress.com/news/2022-12-seizures-faster-decline-people-dementia.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.