

Study considers social determinants of health

December 12 2022



Credit: Pixabay/CC0 Public Domain

Patients with social needs experience a higher number of hospitalizations, obesity, prediabetes and Type 2 diabetes, finds a new study from the Brown School at Washington University in St. Louis on the social determinants of health.

Within the study, patients with [social needs](#) also were disproportionately older, female, Black, uninsured, and living in low-income and high-unemployment neighborhoods, compared with those without what's known as a "Z-code," which are symptoms that are not tied to a specific disorder but still need treatment.

The study found that medical providers faced barriers at multiple levels in documenting social determinants of health.

The paper, "The adoption of [social determinants](#) of [health](#) documentation in clinical settings," was published in the print edition of the journal *Health Services Research* earlier this year. The lead author is Maura Kepper, a research assistant professor at the Brown School at Washington University in St. Louis.

More information: Maura M. Kepper et al, The adoption of social determinants of health documentation in clinical settings, *Health Services Research* (2022). [DOI: 10.1111/1475-6773.14039](https://doi.org/10.1111/1475-6773.14039)

Provided by Washington University in St. Louis

Citation: Study considers social determinants of health (2022, December 12) retrieved 19 April 2024 from <https://medicalxpress.com/news/2022-12-social-health.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--