

Spontaneous baby movements are important for development of coordinated sensorimotor system

December 26 2022





The markers for the motion capture camera were gently applied to the baby's limbs and head and belly, enabling the team to capture the full range of movement. Credit: 2022 Kanazawa et al.

Spontaneous, random baby movements aid development of their sensorimotor system, according to new research led by the University of Tokyo. Detailed motion capture of newborns and infants was combined with a musculoskeletal computer model to enable researchers to analyze communication among muscles and sensation across the whole body.

Researchers found patterns of <u>muscle</u> interaction developing based on the babies' random exploratory behavior that would later enable them to perform sequential movements as infants. Better understanding how our sensorimotor system develops could help us gain insight into the origin of human movement as well as earlier diagnoses of developmental disorders.

Right from birth—and even in the womb—babies start to kick, wiggle and move seemingly without aim or external stimulation. These are called "spontaneous movements," and researchers believe that they have an important role to play in the <u>development</u> of the sensorimotor system, i.e., the ability to control the muscles, movement and coordination. If researchers can better understand these seemingly random movements and how they are involved in early human development, we might also be able to identify early indicators for certain developmental disorders, such as cerebral palsy.

Currently, there is limited knowledge about how newborns and infants learn to move. "Previous research into sensorimotor development has



focused on kinematic properties, muscle activities which cause movement in a joint or a part of the body," said Project Assistant Professor Hoshinori Kanazawa from the Graduate School of Information Science and Technology.

"However, our study focused on muscle activity and sensory input signals for the whole body. By combining a musculoskeletal model and neuroscientific method, we found that spontaneous movements, which seem to have no explicit task or purpose, contribute to coordinated sensorimotor development."

First, the team recorded the joint movements of 12 healthy newborns (less than 10 days old) and 10 young infants (about three months old) using motion capture technology. Next, they estimated the babies' muscle activity and sensory input signals with the aid of a whole-body, infant-scale musculoskeletal computer model which they had created. Finally, they used computer algorithms to analyze the spatiotemporal (both space and time) features of the interaction between the input signals and muscle activity.

"We were surprised that during spontaneous movement, infants' movements 'wandered' and they pursued various sensorimotor interactions. We named this phenomenon 'sensorimotor wandering,'" said Kanazawa. "It has been commonly assumed that sensorimotor system development generally depends on the occurrence of repeated sensorimotor interactions, meaning the more you do the same action the more likely you are to learn and remember it.

"However, our results implied that infants develop their own sensorimotor system based on explorational behavior or curiosity, so they are not just repeating the same action but a variety of actions. In addition to this, our findings provide a conceptual linkage between early spontaneous movements and spontaneous neuronal activity."





This simulation was based on an adult model and infant skeleton. Credit: 2022 Kanazawa et al.

Previous studies on humans and animals have shown that motor behavior (<u>movement</u>) involves a small set of primitive muscular control patterns. These are patterns that can typically be seen in task-specific or cyclic movements, like walking or reaching. The results of this latest study supports the theory that newborns and <u>infants</u> can acquire sensorimotor



modules, i.e., synchronized muscle activities and sensory inputs, through spontaneous whole-body movements without an explicit purpose or task.

Even through sensorimotor wandering, the babies showed an increase in coordinated whole-body movements and in anticipatory movements. The movements performed by the infant group showed more common patterns and sequential movements, compared to the random movements of the newborn group.

Next, Kanazawa wants to look at how <u>sensorimotor</u> wandering affects later development, such as walking and reaching, along with more complex behaviors and higher cognitive functions. "My original background is in infant rehabilitation. My big goal through my research is to understand the underlying mechanisms of early motor development and to find knowledge that will help to promote baby development."

The work is published in the journal *Proceedings of the National Academy of Sciences*.

More information: Kanazawa, Hoshinori, Open-ended movements structure sensorimotor information in early human development, *Proceedings of the National Academy of Sciences* (2022). DOI: 10.1073/pnas.2209953120. doi.org/10.1073/pnas.2209953120

Provided by University of Tokyo

Citation: Spontaneous baby movements are important for development of coordinated sensorimotor system (2022, December 26) retrieved 3 May 2024 from https://medicalxpress.com/news/2022-12-spontaneous-baby-movements-important-sensorimotor.html



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.