

Women and men mistakenly given different advice to prevent heart disease

December 5 2022



Credit: Pixabay/CC0 Public Domain

Women are told to improve their lifestyle to prevent heart disease while men are advised to take statins. That's the finding of a study presented December 3 at [ESC Asia](#), a scientific congress organized by the European Society of Cardiology (ESC), the Asian Pacific Society of Cardiology (APSC), and the Asean Federation of Cardiology (AFC).

"Our study found that women are advised to lose weight, exercise and improve their diet to avoid cardiovascular disease but men are prescribed lipid-lowering medication," said study author Dr. Prima Wulandari of Harvard Medical School and Massachusetts General Hospital, Boston, U.S. "This is despite the fact that guideline recommendations to prevent [heart disease](#) are the same for men and women."

Previous studies have shown that women with cardiovascular disease are given less aggressive treatment compared with men. This study investigated whether these sex differences extend to the prevention of cardiovascular disease.

The study used data from the U.S. National Health and Nutrition Examination Survey (NHANES) from 2017 to 2020. Of 8,512 men and women aged 40 to 79 years and no history of cardiovascular disease, 2,924 participants were at increased risk for developing cardiovascular disease according to a validated risk calculator and therefore eligible to receive statins.

For the 2,924 participants, the researchers calculated the odds of men, compared with women, being prescribed statin therapy, and receiving advice to lose weight, exercise, reduce salt intake, and reduce fat or

calorie consumption. The analyses were adjusted for age, risk of cardiovascular disease, [body mass index](#), resting [heart rate](#), depression score, and education status.

The analysis showed that men were 20% more likely to be prescribed statins compared with women. Compared with men, women were 27% more likely to be advised to lose weight, and 38% more likely to receive recommendations to exercise. Regarding diet, women were 27% more often than men advised to reduce their salt intake, and 11% more frequently told to reduce their fat or calorie consumption.

Dr. Wulandari said, "Following our analysis, we conducted a review of the literature to find possible explanations for the results. This demonstrated that a potential root of the discrepancy in advice is the misconception that women have a lower risk of cardiovascular disease than men. Our findings highlight the need for greater awareness among [health professionals](#) to ensure that both women and men receive the most up-to-date information on how to maintain heart health."

ESC cardiovascular prevention guidelines recommend that adults of all ages should do at least 150 to 300 minutes of moderate intensity, or 75 to 150 minutes of vigorous intensity, aerobic physical activity a week, or an equivalent combination. Everyone should quit smoking. A [healthy diet](#) is recommended, emphasizing plant-based foods including whole grains, fruits, vegetables, pulses, and nuts. Salt should be limited to less than 5 g per day. It is advised that overweight and obese people lose weight to lower [blood pressure](#), [blood lipids](#), and the risk of diabetes, and thereby reduce the likelihood of [cardiovascular disease](#). Statins are recommended according to individual characteristics including age and risk of developing heart disease.

Provided by European Society of Cardiology

Citation: Women and men mistakenly given different advice to prevent heart disease (2022, December 5) retrieved 8 May 2024 from <https://medicalxpress.com/news/2022-12-women-men-mistakenly-advice-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.