

Study suggests the youngest children in class are being overmedicated for ADHD

December 7 2022, by Svein Inge Meland



Credit: Pixabay/CC0 Public Domain

Christine Strand Bachmann has led a study that includes all Norwegian children born between 1989 and 1998, a total of 488,000 people.

The NTNU Ph.D. candidate has reviewed all the prescriptions the children received from the age of 10 until they were 23. She compared children born in January or February with those born in November or December.

"We found major differences in the prescription of ADHD [medication](#). Children born in November-December had an 80% higher risk of being prescribed ADHD medication, compared to those born in January-February. This finding applies to children who were born at term," says Bachmann.

Overmedication

The researcher believes this indicates that the youngest children are being overmedicated.

"I think the increased ADHD medication for the youngest children in the cohort has to do with the way we organize our education system. Put simply, it looks like we're medicating the most immature children because we're comparing them to their oldest classmates, who are a whole year older," she says.

"This shouldn't be the basis for receiving an ADHD diagnosis. We have to expect that a child born in November is going to be less mature than a February child born the same year. Of course we can't stop diagnosing ADHD and giving medicine to those who need it. But what we see here is something else."

Double burden

In particular, Bachmann studied children who were born prematurely, before the 37th week of pregnancy. The use of ADHD medication is

higher for these children than their peers who were born at term. The tendency for the premature children is the same as for full-term—ADHD medication occurs most frequently among the youngest in the class.

"In that sense, the youngest premature children carry a double burden. The disadvantage of being born late in the year comes on top of the disadvantages of being a preemie," says the researcher.

In addition to being a Ph.D. candidate at NTNU's Department of Public Health and Nursing, Bachmann is a pediatrician at St. Olavs Hospital in Trondheim. Her study has been published in the American research journal *Pediatrics*.

Medicating continued longer

Bachmann found another [negative effect](#) for premature children. ADHD prescriptions decreased with age among the full-term children, but the same did not happen for the preterm children who continued to be prescribed ADHD medication at the same level into early adulthood.

"It would appear that children who were born prematurely are extra susceptible to the persistent negative effects of being the youngest in the [school environment](#). Perhaps they experience falling short in the classroom, in gym and in social contexts, compared to their peers. And perhaps the negative experiences settle in the body in a different way and to a greater extent in those born prematurely," she said.

No 'seasonality'

ADHD is characterized by concentration difficulties, hyperactivity and impulsivity beyond what is normal for the child's age.

Some have wondered whether the higher medication use might be because children born in autumn are simply more hyperactive, impulsive and have greater concentration difficulties.

However, Bachmann notes research has not shown any such "seasonal variation."

"We have little reason to believe that the upbringing is different in families with children born late in the year, or that families with better childrearing conditions consciously plan to have children early in the year," she said.

"The findings indicate that the [school](#) situation is the cause of excess ADHD diagnoses. One solution could be for the children in question to wait to start school until the following year," she said.

Flexible school start

Bachmann points out that studies from Denmark, which has a more flexible approach to delayed school starts, do not show an increased use of ADHD medication among the youngest pupils in the class. Forty percent of Danish children born in October-December postpone their school start until the next year.

"In Denmark, parents and kindergartens are involved in assessing a child's school readiness. I believe we could do something similar for our [youngest children](#) in Norway. Premature babies born late in the year could especially benefit from this approach. Some of them would actually have started a year later if they had been full-term."

Only for some

Bachmann is not advocating that all "autumn children" should postpone their school start, which would merely shift the problem.

"We think that this could be relevant for the most vulnerable children. They should be individually assessed, with a focus on the child's particular needs. Having the right follow-up and good support measures for this group is also important, both in the education system and in the healthcare system," she says.

Some would worry that delaying a child's school start would stigmatize them.

"It's difficult to rule that out completely," Bachmann said, "but I still think that it's less problematic than basing our diagnoses on the wrong criteria. That can be just as, if not more, stigmatizing."

Other medicines

The study also looked at the use of other types of medication, such as for depression, anxiety, psychoses and sleeping pills.

"We found the same age effect for several of these drugs as well. Clearly more children born late in the year are receiving this kind of medication, compared to the oldest ones in the class. This was the case for both full-term and preterm babies," Bachmann said.

"But later in adolescence, from ages 15–23, this inequality disappeared. Children who were born late in the year in this age group didn't use more of these medicines than those who were born early in the same year. This also applied to the preemies," says Bachmann.

Data for the study was taken from the Medical birth registry of Norway, the Norwegian Prescription Database, and Statistics Norway.

Norway's Prematurity Association wants a flexible start

Hege Andersson Nordhus, who heads the daily operations of Norway's Prematurity Association (Prematurforeningen), believes the study shows that there is a need for a flexible start to school, especially for preterm babies.

"We've stated on several occasions that premature children, especially those born before what would be their full-term year, should have their school start postponed. It should absolutely be an alternative for children who clearly are not ready for school," says Nordhus.

"At the same time, I believe the school reform from 1997, which introduced Norway's compulsory school start for six-year-olds, should be re-evaluated. It seems that it's difficult to incorporate play into schools as was intended for children in first grade."

Nordhus finds it discouraging to think that the school system is causing unnecessary diagnoses to be inflicted on children.

"We can't prevent children from being themselves just because the child doesn't fit into the school system. It's time to look at the school system and find out what is making this happen," says Nordhus.

More information: Christine Strand Bachmann et al, Relative Age and Psychotropic Drug Use in Preterm and Term-Born Children and Young Adults, *Pediatrics* (2022). [DOI: 10.1542/peds.2022-057085](https://doi.org/10.1542/peds.2022-057085)

Provided by Norwegian University of Science and Technology

Citation: Study suggests the youngest children in class are being overmedicated for ADHD (2022, December 7) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2022-12-youngest-children-class-overmedicated-adhd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.