

Researchers create AI to predict ICU delirium risk

January 26 2023



Credit: Unsplash/CC0 Public Domain

More than one-third of all people admitted to the hospital, and as many as 80% of all patients in an intensive care unit (ICU), develop delirium, a type of brain dysfunction marked by sudden bouts of confusion,

inattention, paranoia, or even agitation and hallucinations. An intensivist at Johns Hopkins Medicine, in collaboration with Johns Hopkins University engineering students, report they have developed artificial intelligence (AI) algorithms that can detect the early warning signs of delirium and can predict—at any time during an ICU stay—a high risk of delirium for a significant number of patients.

"Being able to differentiate between patients at low and high risk of delirium is incredibly important in the ICU because it enables us to devote more resources toward interventions in the high-risk population," says Robert Stevens, M.D., associate professor of anesthesiology and [critical care medicine](#) at the Johns Hopkins University School of Medicine and senior author of the new study results, published in the Dec. 20 issue of the journal *Anesthesiology*. Stevens is also director of precision medicine and informatics, and co-director of the Johns Hopkins Precision Medicine Center of Excellence in Neurocritical Care.

Clinicians already knew that ICU delirium occurs more often in older and sicker patients, and that ICU patients who develop delirium are at higher risk of prolonged hospitalization, future dementia and death. Anti-delirium interventions such as care bundles, changing medications and earlier-than-usual occupational and [physical therapy](#) are effective, experts say, but limited time and resources, as well as the often unpredictable needs of ICU patients, prevent most ICUs from using them in every patient.

The new AI program, designed by undergraduate and master's level engineering students in a precision medicine class taught by Stevens, applied AI algorithms to a publicly available dataset covering more than 200,000 ICU stays at 208 hospitals around the country.

"The underlying idea was that this routinely collected data stored in patients' electronic health records contains signatures that are associated

with delirium risk," says Kirby Gong, a recent master's degree graduate from the Johns Hopkins Department of Biomedical Engineering and first author of the new work.

Using the data, the team developed two computerized models to predict delirium risk. One, a so-called static model, takes a single snapshot of patient data shortly after admission—information about age, severity of illness, other diagnoses, physiologic variables and current medications—to predict delirium risk at any point during a hospital stay. The second, a so-called dynamic model, monitors information over hours and days, including repeat blood pressure, pulse and temperature readings—to provide the patient's continuously updated risk of delirium over the coming 12 hours.

Once the researchers developed the AI models, they tested them on two other sets of data from a Boston hospital, collectively covering more than 100,000 ICU stays. The area under the receiver operating characteristic curve (95% CI) for the first 24-hour model was 0.785, meaning that it was able to predict which patients would get delirium 78.5% of the time. The dynamic model performed even better, predicting [delirium](#)-prone patients up to 90% of the time.

Stevens says he is now testing the models on historical patient data from Johns Hopkins Medicine ICUs, and plans to design a clinical trial to test the use of the algorithms—and how they could shape clinical care—in [patients](#) newly admitted to an ICU. His lab is also applying similar artificial intelligence approaches—often in collaboration with engineering students and faculty—to predict stroke, [heart failure](#), pulmonary embolisms and other emergent events seen in critical care medicine.

"For a lot of these physiological transitions, we think that there are early warning signs that may not be obvious to a clinician but can be picked up

on using the kinds of [artificial intelligence](#)-supported pattern analysis that we used here," says Stevens.

More information: Kirby D. Gong et al, Predicting Intensive Care Delirium with Machine Learning: Model Development and External Validation, *Anesthesiology* (2022). [DOI: 10.1097/ALN.0000000000004478](#).
pubmed.ncbi.nlm.nih.gov/36538354/

Provided by Johns Hopkins University School of Medicine

Citation: Researchers create AI to predict ICU delirium risk (2023, January 26) retrieved 2 May 2024 from <https://medicalxpress.com/news/2023-01-ai-icu-delirium.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.