

COVID-19 drug trial results released

January 4 2023



Credit: Unsplash/CC0 Public Domain

Researchers from the University of Southampton, University of Oxford and Cardiff University have today released findings from a clinical trial investigating the effectiveness of the antiviral treatment molnupiravir against COVID-19—the first treatment tested in the ongoing PANORAMIC trial.

In [their paper](#) published in *The Lancet*, they reported that molnupiravir did not reduce hospitalizations or deaths among higher risk, vaccinated adults with COVID-19 in the community.

The [treatment](#) was, however, associated with a faster recovery time and reduced viral detection and load—participants who received molnupiravir reported feeling better compared to those who received usual care.

Paul Little, Professor of Primary Care Research at the University of Southampton and co-Chief Investigator of PANORAMIC, said, "PANORAMIC is one of the most remarkable studies to have come from [primary care](#), not only in highly innovative recruitment methods during an ongoing pandemic, but in providing timely, clear evidence for patients, their doctors and policy makers for a new antiviral drug—demonstrating that molnupiravir is effective in reducing viral load and helping symptoms, but with no benefit in reducing hospital admissions in a largely vaccinated population."

PANORAMIC Trial

Molnupiravir (brand name Lagevrio) was the first treatment to be studied by the Platform Adaptive trial of NOvel antiViRals for eArly treatMent of COVID-19 In the Community (PANORAMIC), set up to identify which groups of higher risk people were most likely to benefit from new antiviral treatments for COVID-19. The study allows multiple antiviral drugs to be tested in parallel.

Chris Butler, Professor of Primary Care in the Nuffield Department of Primary Care Health Sciences and co-Chief Investigator of PANORAMIC, said, "Finding effective, safe and scalable early treatments for COVID-19 in the community is the next major frontier in our research response to the ongoing worldwide pandemic."

"It is in the community where treatments could have a massive reach and impact. But decisions about who to treat should always be based on evidence from rigorous clinical trials that involve people who would most likely be prescribed the drugs."

He continued: "The evidence PANORAMIC has produced about molnupiravir will guide treatment decisions for COVID-19 world-wide. It is rapidly generating critically important clinical evidence from within the pandemic to guide care during the pandemic itself, in this case determining effects of molnupiravir among people who are almost all vaccinated.

"We must not forget the other ongoing pandemic of [antibiotic resistance](#), which in part stems from using antimicrobial drugs at scale before we did rigorous clinical trials to find out who really benefits from treatment, and who does not. The PANORAMIC team is also doing the necessary trials and gathering evidence about these treatments before we go straight to widespread use."

Professor Richard Hobbs, Head of Oxford Primary Care and co-trial lead, said, "These key findings from 25,000 people in this first PANORAMIC outcome show that the majority of patient groups at moderate to high risk of worse COVID outcomes gained only symptomatic improvement with molnupiravir.

"The low overall rates of serious outcomes reinforces the ongoing importance of the COVID vaccination program in reducing death and hospitalization, with no extra benefit with molnupiravir."

Findings

Study participants were within five days of symptoms onset and either aged over 50 years in [good health](#) or between 18–50 with underlying

health conditions that made them clinically more vulnerable.

A total of 25,786 study participants were randomly assigned to receive either molnupiravir or the usual standard of NHS care.

50 (0.4%) participants experienced serious adverse events in the molnupiravir group and 45 (0.4%) in usual care.

Patients in a sub-group treated with molnupiravir showed reduced viral detection and viral load on Day 7.

Recruitment of people from ethnically and culturally [diverse backgrounds](#) into clinical trials has been important in ensuring audiences likely to benefit from treatment are represented.

Prof Nick Lemoine, Medical Director of the NIHR Clinical Research Network, said, "PANORAMIC is an important NIHR-funded trial that enabled the rapid production of evidence and information about how well this new antiviral works.

"The commendable pace and scale at which this study recruited in at-risk groups shows how we can adapt to make future trials more accessible to a more diverse group of people. It is also testament to the GPs and researchers involved in this trial that we were able to recruit so fast in such record numbers."

The study will now continue to investigate new antiviral medications such as Paxlovid. For more information or to enroll, please visit www.panoramictrial.org.

More information: Christopher C Butler et al, Molnupiravir plus usual care versus usual care alone as early treatment for adults with COVID-19 at increased risk of adverse outcomes (PANORAMIC): an open-label,

platform-adaptive randomised controlled trial, *The Lancet* (2022). [DOI: 10.1016/S0140-6736\(22\)02597-1](https://doi.org/10.1016/S0140-6736(22)02597-1)

Provided by University of Southampton

Citation: COVID-19 drug trial results released (2023, January 4) retrieved 10 May 2024 from <https://medicalxpress.com/news/2023-01-covid-drug-trial-results.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.