

# New study links hearing loss with dementia in older adults

January 10 2023

---



Credit: Pixabay/CC0 Public Domain

A new study led by researchers at the Johns Hopkins Bloomberg School of Public Health found that older adults with greater severity of hearing loss were more likely to have dementia, but the likelihood of dementia

was lower among hearing aid users compared to non-users.

The findings, from a nationally representative sample of more than 2,400 [older adults](#), are consistent with prior studies showing that [hearing loss](#) might be a contributing factor to [dementia risk](#) over time, and that treating hearing loss may lower [dementia](#) risk.

The findings are highlighted in a research letter published online January 10 in the *Journal of the American Medical Association*.

"This study refines what we've observed about the link between hearing loss and dementia, and builds support for [public health](#) action to improve hearing care access," says lead author Alison Huang, Ph.D., MPH, a senior research associate in the Bloomberg School's Department of Epidemiology and at the Cochlear Center for Hearing and Public Health, also at the Bloomberg School.

Hearing loss is a critical public health issue affecting two-thirds of Americans over 70. The growing understanding that hearing loss might be linked to the risk of dementia, which impacts millions, and other adverse outcomes has called attention to implementing possible strategies to treat hearing loss.

For the new study, Huang and colleagues analyzed a nationally representative dataset from the National Health and Aging Trends Study (NHATS). The NHATS has been ongoing since 2011, and uses a nationwide sample of Medicare beneficiaries over age 65, with a focus on the 90-and-over group as well as Black individuals.

The analysis covered 2,413 individuals, about half of whom were over 80 and showed a clear association between severity of hearing loss and dementia. Prevalence of dementia among the participants with moderate/severe hearing loss was 61 percent higher than prevalence

among participants who had normal hearing. Hearing aid use was associated with a 32 percent lower prevalence of dementia in the 853 participants who had moderate/severe hearing loss.

The authors note that many past studies were limited in that they relied on in-clinic data collection, leaving out vulnerable populations that did not have the means or capacity to get to a clinic. For their study, the researchers collected data from participants through in-home testing and interviews.

How hearing loss is linked to dementia isn't yet clear, and studies point to several possible mechanisms. Huang's research adds to a body of work by the Cochlear Center for Hearing and Public Health examining the relationship between hearing loss and dementia.

The study authors expect to have a fuller picture of the effect of hearing loss treatment on cognition and dementia from their Aging and Cognitive Health Evaluation in Elders (ACHIEVE) Study. Results from the three-year randomized trial are expected this year.

**More information:** Alison R. Huang et al, Hearing Loss and Dementia Prevalence in Older Adults in the US, *JAMA* (2023). [DOI: 10.1001/jama.2022.20954](https://doi.org/10.1001/jama.2022.20954)

Provided by Johns Hopkins University Bloomberg School of Public Health

Citation: New study links hearing loss with dementia in older adults (2023, January 10) retrieved 27 April 2024 from <https://medicalxpress.com/news/2023-01-links-loss-dementia-older-adults.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.