


# Study points to practice-changing care for patients with refractory metastatic colorectal cancer

January 23 2023

---



Credit: Unsplash/CC0 Public Domain

Josep Taberero, Director of the Vall d'Hebron Institute of Oncology (VHIO), presented data from the international phase III SUNLIGHT

study during an Oral Abstract Session at this week's American Society of Clinical Oncology's (ASCO) 20th Annual Gastrointestinal Cancers Symposium, January 19-21, San Francisco, CA (U.S.).

This open-label controlled two-arm, phase III comparison study was designed to validate the efficacy and safety of the orally administered combination of trifluridine/tipiracil plus monoclonal antibody bevacizumab versus trifluridine/tipiracil alone in the third-line treatment of patients with refractory metastatic colorectal cancer (mCRC) who had progressed after two lines of prior therapy.

Around 50% of colorectal cancer patients will ultimately develop metastasis, with a five- year survival rate of only 11%. The prognosis for patients with refractory disease remains poor, with median survival generally between 4-8 months. With limited treatment options available, the standard of care for this patient population currently includes trifluridine/tipiracil or regorafenib as monotherapy, or in combination with other approved agents for some patients.

"There is an urgent, unmet clinical need to identify new and more effective treatment strategies to improve the survival and quality of life of our patients. Representing an important step in this direction, the SUNLIGHT phase III study was designed to evaluate the true clinical value of combining trifluridine/tipiracil with bevacizumab," says Josep Tabernero, first author of this present study and Head of the Vall d'Hebron University Hospital's (HUVH) Medical Oncology Department, Vall d'Hebron Barcelona Hospital Campus.

SUNLIGHT enrolled 492 patients across 103 study locations who had received a maximum of two prior chemotherapy regimens for the treatment of advanced colorectal cancer and had demonstrated progressive disease or intolerance to their last regimen. These patients were randomized (1:1) to receive trifluridine/tipiracil in combination

with bevacizumab, or trifluridine/tipiracil monotherapy.

The primary endpoint of this phase III clinical trial was overall survival, and secondary outcome measures were overall response rate, disease control rate, treatment-emergent adverse events, and quality of life.

## **Potentially practice-changing data**

The SUNLIGHT study investigators, including Elena Élez, a Senior Researcher of VHIO's Gastrointestinal and Endocrine Tumors Group, reported an improved median survival of 3.3 months with trifluridine/tipiracil plus bevacizumab, from 7.5 months with trifluridine/tipiracil monotherapy to 10.8 months with the combination regimen.

"An improved survival rate of over three months in metastatic colorectal cancer is considered as statistically as well as clinically relevant. These present data could therefore promise a new standard of care for patients with refractory disease who have progressed after two prior lines of therapy," observes Tabernero.

Progression-free survival was 2.4 months with trifluridine/tipiracil alone versus 5.6 months combined with bevacizumab. Time to deterioration in global health status was 4.7 months and 8.5 months, respectively. Quality of life was graded according to the ECOG Performance Status Scale. Median time to worsening to a grade 2 or more was 9.3 months with the combination compared with 6.3 months in those patients receiving trifluridine/tipiracil alone.

"In addition to the magnitude of benefit observed in patients who received the combination, adding bevacizumab to trifluridine/tipiracil did not increase the risk of serious adverse or adverse events. Improved survival was observed across all subgroups independently of tumor

sidedness, RAS mutational status, and receipt of prior bevacizumab, indicating that the combination regimen is an option for all clinically relevant subgroups," adds Elena Élez, a Senior Consultant of HUVH's Medical Oncology Department.

SUNLIGHT is the first phase III study to validate a new third-line treatment strategy by showing improved efficacy over an existing active regimen in these patients, with a greater benefit than that observed in other trials, and in a population that included patients with poor prognostic factors such as RAS mutations.

"Our data point to trifluridine/tipiracil plus bevacizumab as a new standard of care. This combination therapy could therefore open a much needed, more effective treatment avenue for patients with refractory [metastatic colorectal cancer](#) who have progressed after two lines of therapy," concludes Josep Tabernero.

**More information:** Conference: [condit.com/asco-medical-exhibi ... WvFB-koaAvaOEALw\\_wcB](#)

Meeting: [meetings.asco.org/2023-asco-gi ... tation=215763#215763](#)

Provided by Vall d'Hebron Institute of Oncology

Citation: Study points to practice-changing care for patients with refractory metastatic colorectal cancer (2023, January 23) retrieved 23 April 2024 from <https://medicalxpress.com/news/2023-01-practice-changing-patients-refractory-metastatic-colorectal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.