

# A prescription for prolonged sitting: A five-minute stroll every half hour

January 12 2023

---



Credit: Unsplash/CC0 Public Domain

Mounting evidence suggests that prolonged sitting—a staple of modern-day life—is hazardous to your health, even if you exercise regularly. Based on these findings, doctors advise all adults to sit less and move

more.

But how often do we need to get up from our chairs? And for how long?

Few studies have compared multiple options to come up with the answer most [office workers](#) want: What is the least amount of activity needed to counteract the health impact of a workday filled with sitting?

Now a study by Columbia University exercise physiologists has an answer: just five minutes of walking every half hour during periods of prolonged sitting can offset some of the most harmful effects.

The study, led by Keith Diaz, Ph.D., associate professor of behavioral medicine at Columbia University Vagelos College of Physicians and Surgeons, was published online in *Medicine & Science in Sports & Exercise*, the journal of the American College of Sports Medicine.

Unlike other studies that test one or two activity options, Diaz's study tested five different exercise "snacks": one minute of walking after every 30 minutes of sitting, one minute after 60 minutes; five minutes every 30; five minutes every 60; and no walking.

"If we hadn't compared multiple options and varied the frequency and duration of the exercise, we would have only been able to provide people with our best guesses of the optimal routine," Diaz says.

Each of the 11 adults who participated in the study came to Diaz's laboratory, where participants sat in an ergonomic chair for eight hours, rising only for their prescribed exercise snack of treadmill walking or a bathroom break. Researchers kept an eye on each participant to ensure they did not over- or under-exercise and periodically measured the participants' [blood pressure](#) and blood sugar (key indicators of cardiovascular health). Participants were allowed to work on a laptop,

read, and use their phones during the sessions and were provided standardized meals.

The optimal amount of movement, the researchers found, was five minutes of walking every 30 minutes. This was the only amount that significantly lowered both blood sugar and blood pressure. In addition, this walking regimen had a dramatic effect on how the participants responded to large meals, reducing blood sugar spikes by 58% compared with sitting all day.

Taking a walking break every 30 minutes for one minute also provided modest benefits for [blood sugar](#) levels throughout the day, while walking every 60 minutes (either for one minute or five minutes) provided no benefit.

All amounts of walking significantly reduced blood pressure by 4 to 5 mmHg compared with sitting all day. "This is a sizeable decrease, comparable to the reduction you would expect from exercising daily for six months," says Diaz.

The researchers also periodically measured participants' levels of mood, fatigue, and cognitive performance during the testing. All walking regimens, except walking one minute every hour, led to significant decreases in fatigue and significant improvements in mood. None of the walking regimens influenced cognition.

"The effects on mood and fatigue are important," Diaz says. "People tend to repeat behaviors that make them feel good and that are enjoyable."

The Columbia researchers are currently testing 25 different doses of walking on [health outcomes](#) and testing a wider variety of people: Participants in the current study were in their 40s, 50s, and 60s, and

most did not have diabetes or high [blood](#) pressure.

"What we know now is that for optimal health, you need to move regularly at work, in addition to a daily [exercise](#) routine," says Diaz.

"While that may sound impractical, our findings show that even small amounts of walking spread through the work day can significantly lower your risk of heart disease and other [chronic illnesses](#)."

The study is titled "Breaking Up Prolonged Sitting to Improve Cardiometabolic Risk: Dose-Response Analysis of a Randomized Cross-Over Trial."

The other contributors are Andrea T. Duran (Columbia), Ciaran P. Friel (Feinstein Institutes of Medical Research, Northwell Health), Maria A. Serafini (Columbia), Ipek Ensari (Columbia), and Ying Kuen Cheung (Columbia).

**More information:** Breaking Up Prolonged Sitting to Improve Cardiometabolic Risk: Dose-Response Analysis of a Randomized Cross-Over Trial, *Medicine & Science in Sports & Exercise* (2023).

Provided by Columbia University Irving Medical Center

Citation: A prescription for prolonged sitting: A five-minute stroll every half hour (2023, January 12) retrieved 25 April 2024 from <https://medicalxpress.com/news/2023-01-prescription-prolonged-five-minute-stroll-hour.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.