

Procrastination is linked to poor health—new study

January 17 2023, by Eva Skillgate, Alexander Rozental and Fred Johansson



Credit: Unsplash/CC0 Public Domain

University students have a lot of freedom but not much structure. This can be bad for habitual procrastinators. Studies have shown that at least half of university students procrastinate to a level that is potentially harmful to their education.



But this may not be the only negative result of putting things off until a later date. Studies have found a link between procrastination and <u>poor</u> <u>health</u>. It is associated with higher levels of stress, unhealthier lifestyles and delays in seeing a doctor about <u>health problems</u>.

However, these studies—by the nature of their design—cannot tell us the direction of the relationship. Does procrastination cause poor physical and <u>mental health</u> because people, say, put off starting a new exercise regime or seeing a doctor about a <u>health</u> problem? Or is it the other way around? Does poor physical health, say, lead people to procrastinate because they don't have the energy to do the task now?

To try to solve this riddle, we conducted a <u>longitudinal study</u>—that is, a study that followed people for a period of time, taking measurements at various points in the study. We recruited 3,525 students from eight universities in and around Stockholm and asked them to complete questionnaires every three months for one year.

Our study, published in *JAMA Network Open*, aimed to investigate if students who procrastinate have a higher risk of poor mental and physical health. Of the 3,525 students we recruited, 2,587 answered the follow-up questionnaire nine months later, where several <u>health</u> <u>outcomes</u> were measured.

To understand how procrastination relates to later health outcomes, students with a greater tendency to procrastinate (as scored on a procrastination scale) at the start of the study were compared with students with a lower tendency. The results showed that higher levels of procrastination were associated with somewhat higher symptoms of depression, anxiety and stress nine months later.

Students with higher levels of procrastination were also more likely to report disabling pain in the shoulders or arms (or both), worse sleep



quality, more loneliness and more financial difficulties. These associations remained even when we took other factors that could affect the association into consideration, such as age, gender, parents' level of education, and previous physical and psychiatric diagnoses.

Although no specific health outcome was strongly associated with procrastination, the results suggest that procrastination may be of importance for a wide range of health outcomes, including mental health problems, disabling pain and an unhealthy lifestyle.

As mentioned above, in earlier studies, participants were only assessed at one point in time, making it hard to know which of the conditions came first: procrastination or <u>poor health</u>. By getting students to answer questionnaires at several time points, we could be sure that high procrastination levels were present before we measured their health.

But it is still possible that other factors not accounted for in our analysis could explain the associations between procrastination and subsequent poor health outcomes. Our results are not proof of cause and effect, but they suggest it more strongly than earlier "cross-sectional" studies.

It can be treated

There is good news for habitual procrastinators. <u>Clinical trials</u> (the gold standard of medical research) have shown that cognitive behavioural therapy is effective at reducing procrastination.

The treatment helps the person overcome <u>procrastination</u> by breaking up <u>long-term goals</u> into short-term goals, managing distractions (such as turning off mobile phones), and staying focused on a task despite experiencing negative emotions.

This requires some effort, so it is not something a person can do while



trying to meet a specific deadline. But even small changes can have a large effect. You can try it yourself. Why not start today by leaving your <u>mobile phone</u> in another room when you need to stay focused on a task.

More information: Fred Johansson et al, Associations Between Procrastination and Subsequent Health Outcomes Among University Students in Sweden, *JAMA Network Open* (2023). DOI: 10.1001/jamanetworkopen.2022.49346

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Procrastination is linked to poor health—new study (2023, January 17) retrieved 24 April 2024 from https://medicalxpress.com/news/2023-01-procrastination-linked-poor-healthnew.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.