


Low-grade inflammation may cause arterial stiffness and preclinical atherosclerosis in otherwise healthy adolescents

February 3 2023



Over a 7-year follow-up period from adolescence to young adulthood, increased low-grade inflammation was causally associated with increased arterial stiffness and carotid intima-media thickness, which are signs of premature vascular damage and subclinical atherosclerosis. Credit: Andrew Agbaje

Early vascular damage and atherosclerosis in adolescents may be caused by low-grade inflammation, a paper published in the *Journal of Applied Physiology* concludes. The study was conducted in collaboration between the University of British Columbia in Canada, the University of Bristol in the U.K., the University of Exeter in the U.K., the University of Illinois in the U.S., and the University of Eastern Finland.

In adults, it is known that low-grade inflammation is a major cause of circulatory, musculoskeletal, gastrointestinal, and nervous system diseases and it can also get worse due to aging and poor lifestyle habits. However, in healthy growing children and adolescents, it remains unclear whether low-grade inflammation contributes to premature vascular damage.

"Contrary to some earlier findings, we now know that premature inflammation-induced vascular damage in adolescents, such as higher arterial stiffness, may cause elevated [blood pressure](#), hypertension, [insulin resistance](#), obesity, and elevated [heart rate](#)," says Andrew Agbaje, a physician and clinical epidemiologist at the University of Eastern Finland.

The current study was conducted among 3,862 adolescents who were 17 years old and followed up for 7 years until 24 years of age. The results revealed that about 1 in 3 males and 1 in 2 females were at risk of higher and worsening inflammation by age 24 years. The researchers found that increased inflammation in males may cause higher carotid wall thickness, which is a sign of preclinical atherosclerosis. On the other hand, increased inflammation in females may cause higher arterial stiffness.

According to the researchers, these sex-specific negative effects of inflammation on different aspects of the blood vessels is a novel finding, and it may help understand the pathogenesis of cardiovascular disease in

[early life](#) as well as the optimal approach for treatment. To their surprise, the researchers observed that being overweight or obese did not influence the negative effect of inflammation on premature vascular damage.

"The question that comes to mind is what could cause inflammation in healthy children and adolescents? Unfortunately, the type of diet a child is exposed to may significantly increase inflammation. For example, fruits, vegetables, and fiber-rich foods have significant anti-inflammatory properties and should be encouraged in the family. However, food rich in saturated and trans fats, salty and deep-fried food, red and processed meat, as well as [sugary drinks](#), pastries, and candies may be associated with premature vascular damage since they can significantly raise [inflammation](#)," Agbaje says.

"These novel findings should strengthen our approach to preventing premature [vascular damage](#) in adolescents. Public health experts, pediatricians, and health policymakers should encourage families to take up healthier diet choices. One approach is to significantly lower the prices of fruits and vegetables through government subsidy and tax returns," Agbaje says.

More information: Andrew O. Agbaje et al, Temporal longitudinal associations of carotid-femoral pulse wave velocity and carotid intima-media thickness with resting heart rate and inflammation in youth, *Journal of Applied Physiology* (2023). [DOI: 10.1152/jappphysiol.00701.2022](#)

Provided by University of Eastern Finland

Citation: Low-grade inflammation may cause arterial stiffness and preclinical atherosclerosis in

otherwise healthy adolescents (2023, February 3) retrieved 24 April 2024 from <https://medicalxpress.com/news/2023-02-low-grade-inflammation-arterial-stiffness-preclinical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.