


Women and men shown to have different risk factors for type 2 diabetes

February 15 2023



(A) Receiver-operating characteristic (ROC) curves of fasting IGFBP-2, adiponectin, IGFBP-1 and BMI in detecting T2D in women and men. Logistic regression models not adjusted for confounders. (B) Receiver-operating characteristic (ROC) curves of fasting IGFBP-2, adiponectin and IGFBP-1 combined, as compared to BMI alone or included, in detecting T2D in women and men. Logistic regression models not adjusted for confounders. Credit: *Frontiers in Endocrinology* (2023). DOI: 10.3389/fendo.2022.1092307

Type 2 diabetes is a common disease, affecting 10% of the adult world population, in which heredity and overweight are known risk factors. In a new study at Karolinska Institutet, researchers show that women and men with a predisposition to diabetes have different risk factors forecasting type 2 diabetes and prediabetes. The results were recently published in the journal *Frontiers in Endocrinology*.

In healthy women, low serum level of the adipose tissue protein adiponectin was an independent strong predictor of type 2 diabetes and prediabetes in the future. In healthy men, instead, low serum level of the liver protein IGFBP-1, was an independent strong predictor of type 2 diabetes and prediabetes. This means that these proteins, which are measures of insulin sensitivity in adipose tissue (adiponectin) and liver and muscle (IGFBP-1), can predict whether one has a high risk of getting type 2 diabetes in 10 years.

A previous [study performed in Shanghai in 2016](#) showed [gender differences](#) in the same direction. In men with prediabetes the risk of future type 2 diabetes was significantly reduced if they increased their physical activity and muscle mass. In contrast, the same study showed that women with prediabetes must avoid increasing waist circumference and [abdominal obesity](#) or reducing large waist circumference to prevent type 2 diabetes.

"Our study explains why this gender difference was seen when it came to preventive lifestyles," says Kerstin Brismar, professor of diabetes research at the Department of Molecular Medicine and Surgery. The proteins we studied in men and women increase with increased muscle mass and [physical activity](#) (IGFBP-1) and with reduced abdominal obesity and calorie restriction (adiponectin), she continues.

It may also explain why regular exercise among abdominal obese women is not enough to reduce the risk of type 2 diabetes.

"We have previously shown that [waist circumference](#) was a strong independent predictor of type 2 diabetes in women and now we were able to show that it was linked to reduced production of adiponectin, a protein / hormone produced in the [adipose tissue](#) to, among other things, protect against cell stress," says Kerstin Brismar.

Other studies in both humans and mice have shown that both high normal levels of adiponectin and high normal levels of the liver protein IGFBP-1 protect against getting type 2 diabetes.

More information: Kerstin Brismar et al, Adiponectin, IGFBP-1 and -2 are independent predictors in forecasting prediabetes and type 2 diabetes, *Frontiers in Endocrinology* (2023). [DOI: 10.3389/fendo.2022.1092307](#)

Provided by Karolinska Institutet

Citation: Women and men shown to have different risk factors for type 2 diabetes (2023, February 15) retrieved 9 April 2024 from <https://medicalxpress.com/news/2023-02-women-men-shown-factors-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.