

Adding antipsychotic med to antidepressant may help older adults with treatment-resistant depression

March 3 2023, by Jim Dryden



Credit: Pixabay/CC0 Public Domain

For older adults with clinical depression that has not responded to standard treatments, adding the drug aripiprazole (brand name Abilify)

to an antidepressant they're already taking is more effective than switching from one antidepressant to another, according to a new multicenter study led by Washington University School of Medicine in St. Louis.

Aripiprazole originally was approved by the FDA in 2002 as a treatment for schizophrenia but also has been used in lower doses as an add-on treatment for clinical [depression](#) in [younger patients](#) who do not respond to antidepressants alone.

The new findings are published March 3 in *The New England Journal of Medicine* and are to be presented that same day by Eric J. Lenze, MD—principal investigator and head of the Department of Psychiatry at Washington University—and colleagues at the annual meeting of the American Association for Geriatric Psychiatry in New Orleans.

Many people with clinical depression don't respond to medications used to treat the condition. Consequently, some doctors switch such patients to different antidepressants in the pursuit of finding one that works, while other physicians may prescribe another class of drugs to see if a combination of medications helps.

Both strategies have been recommended by experts as options for [older adults](#) with [treatment-resistant depression](#). However, the new study was designed to help determine which strategy is most effective. Augmenting an antidepressant with aripiprazole helped 30% of patients with treatment-resistant depression, compared to only 20% who were switched to another solo antidepressant, results of the study show.

"Often, unless a patient responds to the first treatment prescribed for depression, physicians follow a pattern in which they try one treatment after another until they land on an effective medication," said Lenze, the Wallace and Lucille Renard Professor and the study's corresponding

author.

"It would be beneficial to have an evidence-based strategy we can rely on to help patients feel better as quickly as possible. We found that adding aripiprazole led to higher rates of depression remission and greater improvements in psychological well-being—which means how positive and satisfied patients felt—and this is good news. However, even that approach helped only about 30% of people in the study with treatment-resistant depression, underscoring the need to find and develop more effective treatments that can help more people."

Treatment-resistant depression is no more or less common in [older people](#) than younger people, but because it seems to accelerate [cognitive decline](#), identifying more effective ways to treat it is very important.

Lenze, along with colleagues at Columbia University, UCLA, the University of Pittsburgh and the University of Toronto, studied 742 people, ages 60 and older, with treatment-resistant depression, meaning their depression had not responded to at least two different antidepressant medications.

The researchers evaluated strategies commonly used in [clinical practice](#) to help alleviate treatment-resistant depression in older patients and designed the study to have two distinct phases. In the first phase, 619 patients, each of whom was taking an antidepressant such as Prozac, Lexapro or Zoloft, were randomly divided into three groups. In the first group, patients remained on whatever antidepressant drug each already was taking but also received the drug aripiprazole (Abilify).

A second group also continued taking antidepressants but added bupropion (brand names Wellbutrin or Zyban), and a third group tapered off of the antidepressant each had been taking and switched to bupropion entirely.

Over the course of 10 weeks, the participants received biweekly phone calls or in-person visits with study clinicians. At these visits, the medications were adjusted according to the individual patient's response and side effects. The researchers found that the group that experienced the best overall outcomes was the one in which patients continued with their original antidepressants but added aripiprazole.

The researchers also anticipated that some people in the study wouldn't respond to the various treatments, so they added a second phase that included 248 participants. In this phase, patients taking antidepressants such as Prozac, Lexapro and Zoloft were treated with lithium or nortriptyline—medications that were widely used before those other, newer antidepressants were approved more than two decades ago. Rates of alleviating depression in the study's second phase were low, about 15%. And there was no clear winner when augmentation with lithium was compared with switching to nortriptyline.

"Those older drugs also are a bit more complicated to use than newer treatments," Lenze explained. "Lithium, for example, requires blood testing to ensure its safety, and it's recommended that patients taking nortriptyline receive electrocardiograms periodically to monitor the heart's electrical activity. Since neither lithium nor nortriptyline were promising against treatment-resistant depression in older adults, those medications are unlikely to be helpful in most cases."

But even the best treatment strategy—adding aripiprazole to an antidepressant—was not markedly successful for many [older patients](#) with treatment-resistant depression.

"This really highlights a continuing problem in our field," said senior author Jordan F. Karp, MD, professor and chair of the Department of Psychiatry at the University of Arizona College of Medicine—Tucson. "Any given treatment is likely to help only a subset of people, and

ideally, we would like to know, in advance, who is most likely to be helped, but we still don't know how to determine that."

Lenze emphasized that overall, antidepressants are highly helpful for the majority of people suffering from clinical depression. At least half of all people with depression feel much better after they begin taking the first medication they try. And almost half of the remainder not helped by a first drug improve when switched to a second drug, But that leaves a sizeable group with [clinical depression](#) that does not respond to two treatments.

The problem is particularly difficult in older adults, many of whom already are taking several medications for other conditions such as high blood pressure, cardiac issues or diabetes," Lenze said. "So switching to new antidepressants every few weeks or adding other psychiatric drugs can be complicated. In addition, because depression and anxiety in older adults may accelerate cognitive decline, there's an urgency to find more effective treatment strategies.

"There definitely is something that makes depression harder to treat in this population, a population that's only going to keep getting larger as our society gets older," he added.

More information: Eric J. Lenze et al, Antidepressant Augmentation versus Switch in Treatment-Resistant Geriatric Depression, *The New England Journal of Medicine* (2023). [DOI: 10.1056/NEJMoa2204462](https://doi.org/10.1056/NEJMoa2204462). www.nejm.org/doi/full/10.1056/NEJMoa2204462

Provided by Washington University School of Medicine

Citation: Adding antipsychotic med to antidepressant may help older adults with treatment-

resistant depression (2023, March 3) retrieved 20 April 2024 from
<https://medicalxpress.com/news/2023-03-adding-antipsychotic-med-antidepressant-older.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.