

Study: Higher fracture risk after total hip replacement when cementless implant used to treat femoral neck fracture

March 8 2023



Credit: Unsplash/CC0 Public Domain

A study by Hospital for Special Surgery (HSS) and other centers found that total hip replacement performed with a cementless prosthesis for a

femoral neck fracture led to a higher rate of a second fracture and subsequent revision surgery. The research was presented today at the American Academy of Orthopaedic Surgeons (AAOS) Annual Meeting in Las Vegas. The results were also published online in *The Journal of Arthroplasty* in October 2022.

Treatments for a femoral neck fracture range from nonoperative management to total [hip replacement](#). When hip replacement is the best treatment option, it can be performed with or without [bone cement](#) to secure the prosthesis.

"Femoral neck fractures are very common in the elderly and are a major cause of morbidity and mortality in this population," explained Alexander McLawhorn, MD, a hip and knee orthopedic surgeon at HSS and study author.

"Several national registry studies have demonstrated a lower risk of overall revision surgery, and specifically revision for periprosthetic fracture, with cemented femoral fixation. Despite this overwhelming data, many patients in the United States receive cementless fixation, including patients who have sustained femoral neck fractures."

"To our knowledge, this was the first non-registry study comparing outcomes of cemented versus cementless total hip replacement for acute femoral neck fractures examining the influence of various patient factors, including bone quality," explained Michael Kheir, MD, who presented the study at the AAOS meeting and played a leading role in the research during his fellowship at HSS.

"The primary purpose was to compare complication rates between both groups, including periprosthetic fractures, aseptic revision, dislocation and mortality rates." Dr. Kheir is currently an [orthopedic surgeon](#) specializing in hip and knee replacement at the University of Michigan.

The retrospective study analyzed 709 total hip replacement cases (199 cemented, 510 cementless) for femoral neck fractures from 2006 to 2020 at three large academic institutions: Hospital for Special Surgery, Indiana University Health and University Hospitals Cleveland Medical Center. Patient demographics, perioperative characteristics and X-rays were reviewed. The Dorr classification system, widely used to evaluate femoral bone quality, categorized patients with Dorr type A, B or C, with type C indicating the weakest bone.

The average age of study participants was 71 years; 66.9% of patients were women. The prevalence of Dorr type bone was 21.1% type A; 66.3% type B; and 12.6% type C. Patients receiving cemented implants were older, had a lower BMI, were more often women and were more likely to have a pre-existing diagnosis of osteoporosis and Dorr C bone type.

Cementless implant stems had a higher all-cause aseptic femoral revision rate (5.1 versus 0.5%,) and periprosthetic femoral fracture rate (4.3 versus 0%). Bone classification played a major role: Each successive Dorr grade had a higher fracture rate with cementless implants (Dorr A =2.3%; Dorr B = 3.7%; Dorr C = 15.9%). There was no difference in rates of dislocation, revision due to infection or mortality rate between groups. The study found an equal distribution of male and female patients who sustained a fracture.

"While femoral fractures occurred in patients with all types of bone quality, Dorr C bone was particularly prone, with an alarmingly high fracture rate when using cementless stems," Dr. Kheir noted. "All of the fractures requiring a revision surgery occurred in cementless cases, suggesting that cemented stems may minimize this complication, regardless of patient sex or Dorr classification."

More information: Michael M. Kheir et al, The Influence of Dorr

Type and Femoral Fixation on Outcomes Following Total Hip Arthroplasty for Acute Femoral Neck Fractures: A Multicenter Study, *The Journal of Arthroplasty* (2022). [DOI: 10.1016/j.arth.2022.10.028](https://doi.org/10.1016/j.arth.2022.10.028)

Provided by Hospital for Special Surgery

Citation: Study: Higher fracture risk after total hip replacement when cementless implant used to treat femoral neck fracture (2023, March 8) retrieved 2 May 2024 from <https://medicalxpress.com/news/2023-03-higher-fracture-total-hip-cementless.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.