

Review: Multiple ways to address telehealth barriers for stroke survivors

March 28 2023



Graphic Abstract: Strategies to Address Barriers to Telehealth Utilization

Graphical abstract. Credit: *Stroke* (2023). DOI: 10.1161/STROKEAHA.122.039566

While the outpatient management of stroke survivors through telehealth



is prone to multiple barriers, it offers many advantages for addressing health equity in stroke survivors, according to a review from UTHealth Houston.

The review—written by Anjail Sharrief, MD, MPH, first author and associate professor of neurology with McGovern Medical School at UTHealth Houston—was published recently in *Stroke*.

Telehealth has seen rapid expansion into chronic care management over the past several years because of the COVID-19 pandemic, Sharrief said. However, there is limited evidence for the benefits of telehealth in addressing disparities in the chronic care of patients living with <u>stroke</u>.

"As we begin to advance the use of telehealth in stroke and in <u>chronic</u> <u>diseases</u> in general, we must consider the potential advantages and barriers to use in populations at highest risk for disparities," said Sharrief, who is also director of stroke prevention at the UTHealth Houston Institute of Stroke and Cerebrovascular Disease and director of the Stroke Transitions Education and Prevention (STEP) clinical program.

"While telehealth can expand access to care and treatment in many ways, it also has potential to increase disparities in populations with lower levels of digital literacy, limited access to internet, and in whom physical and cognitive limitations pose barriers to telehealth utilization."

In the paper, Sharrief and other researchers listed the various benefits telehealth offers for a range of stroke complications—such as impaired gait, vision, and cognition—as well as for certain social determinants of health, including economic instability, geographic location, and limited social support.

For example, for patients dealing with stroke-related disabilities, the



researchers found that telehealth addresses barriers related to mobility challenges and special equipment needed to access clinic spaces by removing the need for transportation altogether. It also decreases the number of in-person visits required of patients, allowing for better multidisciplinary care and the ability to remotely monitor <u>blood pressure</u> and cardiac arrhythmias.

"One important thing to consider is that while we focus on stroke-related disability, the same principles apply to patients with other neurological diseases," Sharrief said.

The team also identified key barriers stroke survivors face when it comes to telehealth—including <u>physical disabilities</u> from stroke, cognitive disabilities from stroke, limited access to internet, limited digital literacy, and limited English proficiency—and outlined potential strategies to address them.

Their key recommendations for accommodating stroke survivors with limited internet access and limited digital literacy include:

- Utilize <u>social workers</u> or community health workers to connect patients to available federal programs, such as the Affordable Connectivity Program, which offers discounts for internet access as well as discounts toward the purchase of a laptop, desktop computer, or tablet.
- Provide mobile hotspot devices to patients with limited Wi-Fi access.
- Use community-based telehealth outposts for visits.
- Use cellular devices for telemonitoring (e.g., blood pressure monitoring), instead of services or devices requiring Bluetooth.
- Use texting, secure messaging, smart phone applications, and other tools that do not require high-speed internet.
- Develop <u>digital tools</u> to accommodate patients with impaired



literacy or cognitive limitations.

- Provide digital literacy training as a component of interventions.
- Use digital health navigators.
- Include patients with impaired digital literacy in intervention or study design.

"Several of the listed recommendation have shown promise for improving telehealth access and utilization in other chronic disease populations," Sharrief said. "My group and co-authors on the manuscript are testing these in the stroke population."

Additionally, to accommodate patients with hemiparesis or incoordination resulting from stroke, the researchers recommend developing telehealth tools and engaging informal caregivers and family members to participate in telehealth interventions. For <u>stroke survivors</u> with cognitive issues, they suggest avoiding the use of some tools, such as platforms and monitoring equipment, that require multiple steps for setup or need regular troubleshooting.

For patients with limited English proficiency, the authors encourage the development of <u>telehealth</u> platforms, telemonitoring tools, and other digital health tools to accommodate patient and caregiver language preferences.

More information: Anjail Z. Sharrief et al, Telehealth Trials to Address Health Equity in Stroke Survivors, *Stroke* (2023). DOI: 10.1161/STROKEAHA.122.039566

Provided by University of Texas Health Science Center at Houston

Citation: Review: Multiple ways to address telehealth barriers for stroke survivors (2023, March



28) retrieved 6 May 2024 from <u>https://medicalxpress.com/news/2023-03-multiple-ways-telehealth-barriers-survivors.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.