

Sniffing urinary acetone as a fast way to diagnose type 2 diabetes

March 10 2023, by Zhang Nannan



A brief schematic diagram of the detection principle and method. Credit: Xu Wei

Type 2 diabetes (T2D) accounts for more than 90% of the confirmed diabetes cases. It has become a common metabolic disease and is expected to affect 380 million people worldwide by 2025. Currently, clinically used diagnostic tests are mainly based on fasting plasma glucose, oral glucose tolerance test and glycosylated hemoglobin A1c



(HbA1c). Although these methods are highly accurate, they are all invasive tests based on blood sampling.

A research team led by Prof. Shen Chengyin from the Hefei Institutes of Physical Science (HFIPS) of the Chinese Academy of Sciences (CAS) has been searching for a new non-invasive screening method for T2D.

Using a self-developed proton transfer reaction-mass spectrometry (PTR-MS), the researchers detected <u>volatile organic compounds</u> in the urine of T2D patients and the results were published in *Talanta*.

The study was based on the detection of urinary acetone by PTR-MS.

Urinary acetone is one of the ketones produced by fat metabolism. Due to insufficient utilization and storage of glucose in the blood, T2D patients will accelerate the metabolism of urinary acetone.

The researchers recruited 180 T2D patients and 180 healthy volunteers for a multicenter study.

A diagnostic model with an <u>accuracy</u> of 81.3% and a threshold of 690.1 ppbv was established by using urinary acetone at center one. The model was validated at two other centers with similar results. Furthermore, the accuracy of this method was comparable to the <u>diagnostic method</u> used in clinical practice.

"Sniffing T2D by urinary acetone is safe, non-invasive, fast and accurate," said Xu Wei, first author of the study. "We hope that the method can provide a reference for screening and diagnosis of T2D."

More information: Wei Xu et al, Rapid and non-invasive diagnosis of type 2 diabetes through sniffing urinary acetone by a proton transfer reaction mass spectrometry, *Talanta* (2023). <u>DOI:</u>



10.1016/j.talanta.2023.124265

Provided by Chinese Academy of Sciences

Citation: Sniffing urinary acetone as a fast way to diagnose type 2 diabetes (2023, March 10) retrieved 8 May 2024 from https://medicalxpress.com/news/2023-03-sniffing-urinary-acetone-fast-diabetes.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.