

New tool to diagnose and assess the severity of sarcopenia

March 13 2023



Grade 1 (normal muscle) biceps brachii. (A) MSUS short axis view, (B) anatomical image, (C, D) histology. MSUS, musculoskeletal ultrasound. Credit: *RMD Open* (2023). DOI: 10.1136/rmdopen-2022-002779

Sarcopenia is a degenerative disease characterized by a pathological decrease in muscle strength that particularly affects older people. Researchers at the University of Barcelona have developed a new tool to



assess the presence and severity of this muscular deterioration.

The study, published in the journal *RMD Open*, provides a muscle quality scoring system based on musculoskeletal ultrasound scans, which has been validated with anatomical and histological samples from patients. It is the first imaging method that assesses muscle quality, an essential factor in the disease. According to the researchers, these results could be useful "not only for diagnostic purposes but also for patient monitoring in clinical practice and clinical trials."

Researchers from the Unit of Human Anatomy and Embryology and the Histology Unit of the Department of Pathology and Experimental Therapeutics of the Faculty of Medicine and Health Sciences (Bellvitge Campus) of the UB have participated in the study. Experts from the Puigvert Foundation, the University of Genoa, the University of Oslo and the University of Copenhagen also took part in the study.

A harmless and low-cost technique

A key feature of sarcopenia is the deterioration of muscle quantity and quality, which worsens the quality of life and is associated with cardiorespiratory diseases and an increased risk and prolonged hospital admission. A wide range of tests and tools are now available for disease characterization in practice and research.

"In most cases, diagnosis and progression are measured by functional tests or patient surveys on aspects such as falling, feelings of weakness, slowness in walking, difficulty getting up from a chair or <u>weight loss</u>, which provide information on the patient's condition," notes Ingrid Möller, researcher at the UB and the Poal Institute of Rheumatology (Barcelona) and first author of the article.

In this context, <u>imaging techniques</u> play a critical role in the objective



assessment of sarcopenic patients. However, most of them, such as dualenergy X-ray absorptiometry (DXA), <u>magnetic resonance</u> imaging (MRI) or computerized tomography (CT), focus on muscle quantity. "These techniques show the quantity of muscle, but muscle quality —which is a relatively new term— is more important, as the quantity is very variable due to different factors such as age or height," she says.

Currently, there is no universal consensus on appropriate methods of assessing muscle quality in regular clinical practices. In this situation, the researchers propose the use of musculoskeletal ultrasound, as it is a technique that can show muscle texture and changes in micro- and macroscopic aspects of muscle architecture and composition caused by sarcopenia. Moreover, unlike other imaging techniques, it is a low-cost, portable and harmless test, since it does not emit ionizing radiation.

A method validated with clinical measurements

To confirm the diagnostic validity of this new tool, the study assessed one muscle from the lower extremity (rectus femoris) and one from the upper extremity (biceps brachii) from ten donor cadavers aged 68 to 91. To do this, two researchers assessed and qualitatively scored the severity of muscle degeneration from ultrasound images and then compared the results with the anatomical and histological assessment of the cadavers.

The assessment was performed blind to the other assessor and was repeated half an hour later by each of the researchers. In addition, an assessment by an experienced histologist under the same methodological conditions was also carried out. According to the researchers, this methodology is one of the "strengths of the study, as it allows direct comparison between imaging and clinical measurements of muscle quality."

The next challenge for the researchers is to verify the scoring system in



patients and test the added value of the tool in the long-term monitoring of patients with sarcopenia, as well as its application in clinical trials related to it. In this sense, as the researcher explains, the team from the University of Copenhagen that has participated in this work is going to apply the new tool in a cohort of patients to see how it works in terms of prognostic value.

"We believe it works very well on diagnostics, but both in this case and in its prognostic value, it needs to be corroborated in longitudinal studies with patients," explains Ingrid Möller.

A potential tool for other diseases

Muscle quality is an important factor in other muscle diseases as well as sarcopenia. Thus, for the researchers, musculoskeletal ultrasound could also have applications in other disorders or even in the rehabilitation of sportsmen and women, as it will allow them to gain a deeper understanding of <u>muscle</u> quality, its variations in different immunological, degenerative, and metabolic diseases, and the monitoring of treatments.

"It is the starting point in a field in which there is no similar test, so we believe that it is a tool that will be extended to this and other disease," concludes the researcher.

More information: Ingrid Möller Parera et al, Ultrasound assessment of degenerative muscle sarcopenia: the University of Barcelona ultrasound scoring system for sarcopenia, *RMD Open* (2023). DOI: 10.1136/rmdopen-2022-002779

Provided by University of Barcelona



Citation: New tool to diagnose and assess the severity of sarcopenia (2023, March 13) retrieved 23 June 2024 from <u>https://medicalxpress.com/news/2023-03-tool-severity-sarcopenia.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.