

## **3D** heart modelling offers non-invasive diagnosis and treatment options for aortic stenosis

April 4 2023



Credit: King's College London

The 3D printing of intricate anatomical structures has become increasingly sophisticated in line with advances in imaging technology, with the use of printed phantoms facilitating the more detailed manipulation of a simulated valve's anatomy and control of blood flow



behavior.

King's researchers from the Cardiac Modeling and Imaging Biomarkers group have published two new papers detailing advances in efforts to deploy 3D printed heart models (phantoms) to simulate and study <u>aortic</u> <u>stenosis</u>.

Aortic stenosis is the condition of calcified and thickened aortic heart valves that obstructs <u>blood flow</u>.

Computer modeling and 3D printing of aortic flow phantoms offer an alternative to in vivo studies which carry both patient recruitment difficulties and potential procedural risks compared to the simulated alternative which can bear higher variations in <u>blood</u> pressure flow and drop.

The research published in the *Journal of Cardiovascular Magnetic Resonance* looked at the development of a non-invasive method to access the pressure of blood flow momentum via <u>cardiovascular magnetic</u> <u>resonance</u> (CMR). The study published in the *Journal of Cardiovascular Translational Research* looked at blood flow dynamics and measurements in phantoms models.

"By developing valve models that behave like real human valves, new techniques which more accurately characterize the severity of disease can be developed and improved without disrupting patients' care," says Harminder Gill, BM BCh.

Current techniques used to measure the severity of aortic stenosis such as Doppler echocardiography can be subject to uncontrolled sources of error and require invasive pressure measurements for the patient. The use of aortic <u>flow</u> phantoms offering one solution to this challenge.



"The decision on how and when to treat stenotic valves is complex and the diagnositc tools typically used in clinical routine have barely evolved during the past 50 years. Thus advances in the study of aortic stenosis patho-physiology are essential to provide a more comprehensive caracterization of this condition. The non-invasive assessment of the pressure recovery distance allows the detection of invasive catheterization errors as well as understanding the vessel length required for haemodynamic homeostasis to be reached," says Joao Filipe Fernandes, Ph.D.

"These advances will enable us to take well informed decision on the best balance between drugs and surgeries for people living with valve conditions," says Prof. Pablo Lamata, head of Cardiac Modeling and Imaging Biomarkers Group.

**More information:** Joao Filipe Fernandes et al, Non-invasive cardiovascular magnetic resonance assessment of pressure recovery distance after aortic valve stenosis, *Journal of Cardiovascular Magnetic Resonance* (2023). DOI: 10.1186/s12968-023-00914-3

Harminder Gill et al, Aortic Stenosis: Haemodynamic Benchmark and Metric Reliability Study, *Journal of Cardiovascular Translational Research* (2023). DOI: 10.1007/s12265-022-10350-w

Provided by King's College London

Citation: 3D heart modelling offers non-invasive diagnosis and treatment options for aortic stenosis (2023, April 4) retrieved 9 May 2024 from <u>https://medicalxpress.com/news/2023-04-3d-heart-non-invasive-diagnosis-treatment.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.