

Adolescents who use smartphones for more than three hours a day found to suffer more from back pain

April 6 2023



Credit: Pixabay/CC0 Public Domain


Thanks to the popularization of smartphones and tablets, and the multiplication of video channels, computer games and educational apps,

children and adolescents are spending more and more time looking at screens, usually with bad posture, which can cause back pain, among other problems.

A study by Brazilian researchers reported in an article published in *Healthcare* identified several risk factors for spinal health, such as looking at screens for more than three hours a day, proximity of the eyes to the screen, and sitting or lying on the stomach.

The study focused on thoracic spine pain (TSP). The thoracic spine is located at the back of the chest (the thorax), mostly between the shoulder blades, extending from the bottom of the neck to the start of the lumbar spine. The data analyzed came from surveys of 14- to 18-year-old male and female students in the first and second years of high school in Bauru, a medium-sized city in São Paulo state.

A baseline questionnaire was completed in March-June 2017 by 1,628 participants, of whom 1,393 completed a follow-up questionnaire in 2018. The analysis showed a one-year prevalence of 38.4% (the proportion reporting TSP in both the baseline and follow-up surveys) and a one-year incidence of 10.1% (new TSP reported only in the follow-up survey). More girls than boys reported TSP.



Flow chart of study participants (TSP, thoracic spine pain). Credit: *Healthcare* (2023). DOI: 10.3390/healthcare11020196

Risk factors

TSP is common in different age groups of the general population worldwide, with prevalence ranging from 15%-35% in adults and 13%-35% in children and adolescents. Explosive growth in the use of electronic devices during the COVID-19 pandemic clearly made the problem worse.

The risk factors associated with TSP are physical, physiological, psychological and behavioral, according to several investigations. There is also strong evidence of the effects of physical activity, sedentary

habits and mental disorders on spinal health. All these factors are considered critical by the World Health Organization (WHO) in its latest global review of evidence and guidelines.

"The study can be used to inform [health education](#) programs for school students, teachers, staff and parents," said Alberto de Vitta, first author of the article. He has a Ph.D. in education from the State University of Campinas (UNICAMP) and completed a postdoctoral fellowship in [public health](#) at São Paulo State University (UNESP) in Botucatu.

"This is in line with some of the objectives of the National Curriculum Parameters [PCN, Brazilian government guidelines for [secondary schools](#)], according to which schools are responsible for health education, including identification of risks to individual and collective health and interventions to combat them, as well as promotion of self-care habits with regard to the body's possibilities and limits," said Vitta.

Information on risk factors for TSP in [high school students](#) is important because children and adolescents with back pain are more inactive, achieve less academically and have more psychosocial problems, according to the article. In addition, fewer studies have been conducted on TSP than on lower back and neck pain. A systematic review of the literature on TSP found only two prospective studies regarding prognostic factors.

More information: Alberto De Vitta et al, Thoracic Spine Pain in High School Adolescents: A One-Year Longitudinal Study, *Healthcare* (2023). [DOI: 10.3390/healthcare11020196](https://doi.org/10.3390/healthcare11020196)

Provided by FAPESP

Citation: Adolescents who use smartphones for more than three hours a day found to suffer more from back pain (2023, April 6) retrieved 11 May 2024 from

<https://medicalxpress.com/news/2023-04-adolescents-smartphones-hours-day-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.