

Matching form and function of brain cell types

April 25 2023



Extracellular action potential (EAP) recordings from in vivo epxeriments and biophysically realistic single-cell models. Extracellular action potential (EAP) recordings from in vivo experiments (**a**–**d**) and single-cell modeling (**e**–**g**). **a** Left, labels for Cre-line and morphology (spiny vs. aspiny) groups of single neurons used in this study characterized in vitro via intracellular electrophysiology and morphology reconstructions. Right, in vivo EAP waveform analysis typically results in two clusters, fast-spiking (FS) vs. regularspiking (RS) units. **b** Primary visual cortex (V1) in the mouse brain (left) and



typical cortical depth placement of a Neuropixels probe along V1 (right, from the Allen Reference Atlas-Mouse Brain⁸⁶). **c** The 384 electrode sites of the Neuropixels probe are densely arranged along the linear shank probe (left; 20 µm vertical spacing, 2 sites per row; black squares: location of recording sites). EAP waveforms from two example units (unit #1: FS; unit #2: RS), including the channel with the largest amplitude (closest to the soma, bolded lines) and channels above and below the soma. d Top: the number of Neuropixelsimplanted mice for wild-type (n = 24), parvalbumin-expressing (Pvalb, n = 8), and somatostatin-expressing (Sst, n = 12); bottom: the distribution of units per wild-type mouse recorded in V1 during drifting gratings (total number of units = 204). Distribution of units along the V1 depth axis with 0 indicating the center of layer 4. e Bio-realistic, single-cell models of V1 ("all-active") are generated from in vitro experiments and activated via synaptic background to elicit intracellular activity and associated EAP signals in the vicinity of the cellular morphology. The cellular morphology is represented with a spherical soma and full dendritic reconstruction (axon not shown). Example simulations of EAP signals are shown for a spiny (top: red, cell ID: 395830185) and an aspiny (bottom: blue, cell ID: 469610831) single-cell model. f Four examples of the multi-channel EAP, including the channel with the largest amplitude (bolded lines, closest to the soma) and channels above and below the soma (top: 2 spiny models; bottom: 2 aspiny models). g In total, 33 single-cell models (15 spiny and 18 aspiny) were generated using a computational optimization framework and included in the study covering a range of major reporter lines and cortical depths. Source data are provided as a Source Data file. Credit: Nature Communications (2023). DOI: 10.1038/s41467-023-37844-8

Investigators at Cedars-Sinai have created computer-generated models to bridge the gap between "test tube" data about neurons and the function of those cells in the living brain. Their study, published in the journal *Nature Communications*, could help in the development of treatments for neurological diseases and disorders that target specific neuron types based on their roles.



"This work allows us to start looking at the <u>brain</u> like the complex machine that it is, rather than as one homogenous piece of tissue," said Costas Anastassiou, Ph.D., a research scientist in the departments of Neurology, Neurosurgery and Biomedical Sciences at Cedars-Sinai and senior author of the study. "Once we are able to distinguish between the different cell types, instead of saying that the entire brain has a disease, we can ask which <u>neuron types</u> are affected by the disease and tailor treatments to those neurons."

Neurons are the main functional units of the brain. The signals passing through these cells—in the form of electrical waves—give rise to all thought, sensation, movement, memory and emotion.

The study used data from <u>laboratory mice</u> to establish a new method for examining relationships between neuron type and function, and focused on the mouse <u>primary visual cortex</u>, which receives and processes <u>visual</u> <u>information</u>. It is one of the best studied parts of the brain—both in vitro, where tissue is studied in a dish or test tube outside the <u>living</u> <u>organism</u>, and in vivo, where it is studied in the living animal.

The investigators' goal was to link the two worlds.

"Based on in vitro studies of genetic makeup and physical structure, we know something about what various classes of neurons look like, but not their function in the living brain," Anastassiou said. "When we record the activity of brain cells in vivo, we can see what neurons are responding to and what their function is, but not which classes of neurons they are."

To link form to function, investigators first used in vitro information to create computational models of various types of neurons and to simulate their signaling patterns.



They next took advantage of the newest technology in single-neuron recording to observe activity in the brains of laboratory mice while the mice were exposed to different sorts of visual stimuli. Based on the shapes of the signals or "spikes" of <u>neurons</u> in response to visual input, investigators separated the cells they recorded into six groups.

"Once we had our models and our in vivo data, the fundamental question was which computational models produced the most similar signaling shape and waveform to each of the six in vivo clusters we identified, and vice versa," Anastassiou said. "Not all of the in vivo clusters and models matched perfectly, but some did."

More data, and possibly experiments involving more-sophisticated visual stimuli, might be required to match all the computational models and cell clusters, and Anastassiou said that future studies will be dedicated to perfecting the method established in the current paper.

"There's a wealth of information about the identity of cell types in the human brain, but not about the role of those cell types in cognitive functioning or how they are affected by disease," Anastassiou said. "Now there is a window through which we can look at these things and ask these questions. It's clear that we have a long way to go, but we're excited about the next steps in this journey."

The ultimate goal is to pave the way for discoveries that change patients' lives.

"Our research scientists are continually striving to expand our knowledge of the workings of the <u>human brain</u> at the most detailed level," said Keith L. Black, MD, chair of the Department of Neurosurgery and the Ruth and Lawrence Harvey Chair in Neuroscience at Cedars-Sinai. "Pinning down the specific type and function of each neuron may one day lead to the discovery of lifesaving treatments for brain diseases and



neurological disorders."

More information: Yina Wei et al, Associations between in vitro, in vivo and in silico cell classes in mouse primary visual cortex, *Nature Communications* (2023). DOI: 10.1038/s41467-023-37844-8

Provided by Cedars-Sinai Medical Center

Citation: Matching form and function of brain cell types (2023, April 25) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2023-04-function-brain-cell.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.