

## Iron found to be an important regulator of neurodifferentiation

April 24 2023



Depletion of IRPs accelerated ROS accumulation. Credit: Science China Press

Iron is important for life, and iron deficiency impairs development, but whether the iron level regulates neural differentiation remains elusive. Iron-regulatory proteins (IRPs), including IRP1 and IRP2, are the major regulators of cellular iron homeostasis.



However, as fetal mice with deficiencies of both IRP1 and IRP2 are embryonic lethal, it has been difficult to explore the role of IRPs in the <u>nervous system</u>, especially in the early stage of neurodevelopment.

In a study published in the journal *Science China Life Sciences*, Shiyang Chang, Peina Wang and their colleagues in Dr. Chang's group and Dr. Wang's group bypassed the bottleneck of embryonic lethality in IRP1<sup>-/-</sup>IRP2<sup>-/-</sup> mice and generated IRP1<sup>-/-</sup>IRP2<sup>-/-</sup> embryonic stem cells (ESCs) to evaluate the physiological significance of IRPs in neurodevelopment.

The team found that the IRPs knockout ESCs showed severe <u>iron</u> <u>deficiency</u>, and meantime the Pax6- and Sox2-positive neuronal precursor cells and Tuj1 fibers in IRP1<sup>-/-</sup>IRP2<sup>-/-</sup> ESCs were significantly decreased during neural differentiation. Consistently, in vivo study showed that the knockdown of IRP1 in IRP2<sup>-/-</sup> fetal mice remarkably affected the differentiation of neuronal precursors and the migration of neurons.

## Medical X press





Iron is an important factor for neurodifferentiation. Credit: Science China Press

Iron supplementation could reduce the effects of IRPs deficiency on neural development, albeit excessive <u>iron</u> inhibited neural development. These findings suggest that iron is a key factor in neurodifferentiation.

"We were absolutely thrilled by the power of iron to control neurodifferentiation so effectively. Most excitingly, when supplementing iron-deficient ESCs with iron, these ESCs could differentiate normally. This may bring new therapies for the treatment of neurological diseases in the future," Yan-Zhong Chang—the lead author of the study noted.

The authors further demonstrated that the underlying mechanism of the impaired neurodifferentiation was in association with an increase in reactive oxygen species (ROS) production caused by the substantially low level of iron and the downregulation of iron-sulfur cluster protein ISCU, which, in turn, affected the proliferation and differentiation of stem cells. Thus, sufficient iron is crucial for maintaining normal neural differentiation, which is termed as ferrodifferentiation.

**More information:** Shiyang Chang et al, Ferrodifferentiation regulates neurodevelopment via ROS generation, *Science China Life Sciences* (2023). DOI: 10.1007/s11427-022-2297-y

Provided by Science China Press

Citation: Iron found to be an important regulator of neurodifferentiation (2023, April 24)



retrieved 1 May 2024 from <u>https://medicalxpress.com/news/2023-04-iron-important-neurodifferentiation.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.