


Research links common insecticide to neurodevelopmental disorders

April 25 2023



NND-related behavioral changes in DPE mice. A) The experimental design, showing pesticide exposures in the dam (3 mg/kg every 3 days) for 2 weeks preconception, then throughout pregnancy and lactation. Time points for testing of offspring are shown. B) DPE male pups produced fewer USVs than controls, while female pups were unaffected by DPE. C) DPE mice (N = 14) showed more repetitive digging than controls (N = 19) in a marble-burying test. D) DPE male mice (N = 14) performed more repetitive self-grooming than controls (N = 19) in a novel cage observation. E) DPE male mice (N = 15) showed no

difference from controls (N = 18) on within-session fear acquisition. F) Following classic fear conditioning, DPE male mice (N = 15) showed less freezing than controls (N = 18) in the fear recall test. G) DPE male mice (N = 15) were less likely than controls (N = 17) to acquire the operant conditioning response to nose poke for food after 7 days of training. Error bars are SEM. *P² (G). Credit: *PNAS Nexus* (2023). DOI: 10.1093/pnasnexus/pgad085

A new study from The University of Toledo suggests early exposure to a common class of insecticides called pyrethroids may increase the risk of autism and other developmental disorders, even at levels currently recognized as safe by federal regulators.

The findings, which come from a study of mice, were published in *PNAS Nexus*.

Pyrethroids are some of the most widely used insecticides in the country, appearing in both consumer products and industrial preparations.

"If you have someone who comes and sprays in your house, this is likely what they're spraying. It's used in landscaping, it's what they fog in the streets for mosquitos. It's everywhere," said Dr. James Burkett, an assistant professor of neuroscience in the UToledo College of Medicine and Life Sciences and the paper's corresponding author. "Our study, however, adds to the evidence that these chemicals might not be as safe for children and pregnant women as we once believed."

Interest in a possible link between autism and pyrethroids has grown after several [epidemiological studies](#) documented higher rates of [neurodevelopmental disorders](#) in areas where the pesticides were used.

The new UToledo-led research sought to build on those population-based

studies by analyzing the specific behavioral changes attributable to low-level exposure to pyrethroids.

Working with a team that included scientists from Columbia, Emory, and the University of Southern California, Burkett examined the offspring of female mice who were exposed to small doses of the [pyrethroid](#) insecticide deltamethrin before, during and immediately after pregnancy.

The researchers found those mice exhibited increased hyperactivity and repetitive behaviors, less vocalization, and were more likely to fail basic learning tests compared to controls.

The young mice also experienced disruptions in their dopamine system.

"These are all similar to symptoms [human patients](#) with neurodevelopmental disorders might have," Burkett said. "We are not saying these mice have autism or that they have ADHD. That's not the goal here. What we are saying is that something in their brain has been altered by this exposure and it's resulting in the same kinds of behaviors that we see in children with autism."

Dr. Gary Miller, vice dean for research strategy and innovation at Columbia University Mailman School of Public Health and a co-author on the study, said the research builds on prior efforts to identify other potentially harmful pest control chemicals.

"We have reduced our exposures to many classes of dangerous pesticides over the past few decades through restrictions and regulations." Miller said. "This study adds to a growing body of literature that the widely used pyrethroids are not without adverse effects and should be further evaluated for their safety."

The prevalence of autism in the United States has risen significantly over the past two decades, rising from one in 150 children in 2000 to one in 44 children in 2018, according to the Centers for Disease Control and Prevention.

Some of that increase is being driven by more awareness of autism spectrum disorder and better diagnostic processes. However, scientists still don't know what causes most cases of autism.

"Research has traditionally focused on genetics, but it's come to be appreciated there's also a significant environmental element," Burkett said. "Unfortunately, those factors are very poorly understood. That's why this kind of research is so important. You can't change someone's genetics, but you can address environment factors."

Pyrethroids are not the only class of pesticides that have been linked to autism, and scientists generally agree that development of the condition requires more than one single trigger.

"This research represents a piece of the puzzle. It is not definitive proof that the pesticide is unsafe or directly causing [autism](#) in humans," Burkett said. "It may, however, suggest that the safe level of the pesticide needs to be revisited for pregnant women and children."

More information: Melissa A Curtis et al, Developmental pyrethroid exposure causes a neurodevelopmental disorder phenotype in mice, *PNAS Nexus* (2023). [DOI: 10.1093/pnasnexus/pgad085](https://doi.org/10.1093/pnasnexus/pgad085). [academic.oup.com/pnasnexus/art ... /2/4/pgad085/7128809](https://academic.oup.com/pnasnexus/article/2/4/pgad085/7128809)

Provided by University of Toledo

Citation: Research links common insecticide to neurodevelopmental disorders (2023, April 25)
retrieved 18 April 2024 from

<https://medicalxpress.com/news/2023-04-links-common-insecticide-neurodevelopmental-disorders.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.