


Nightly variation in sleep apnea sharply increases cardio risks, shows study

April 26 2023



Association between mean apnea–hypopnea index, apnea–hypopnea index variability and hypertension. Marginal probability of hypertension in relation to: **a** mean apnea–hypopnea index and **b** apnea–hypopnea index variability.

Poorly managed sleep apnea can lead to uncontrolled blood pressure—a little known underlying risk for millions of people's cardiovascular health, a new study in *npj Digital Medicine* reveals.

Most at risk are those with wide-ranging variability in their night-to-night Obstructive Sleep Apnea (OSA) severity which can be overlooked

or misdiagnosed during standard clinical testing, the Flinders University-led research warns.

"Regardless of their overall OSA average severity, people with increased night-to-night variability in the amount of [sleep apnea](#) are at a 50% to 70% increased likelihood of having uncontrolled hypertension and variable blood pressure—a confirmed risk factor for cardiovascular events, all-cause mortality, vascular organ damage, [atrial fibrillation](#) and dementia," says Dr. Bastien Lechat, lead author in the new article.

"High variability in OSA severity is an independent clinical predictor of [uncontrolled hypertension](#) and has significant implications for identifying patients most at risk of cardiovascular harm."

OSA is a common clinical sleep disorder characterized by repetitive upper airway collapse during sleep and is estimated to affect approximately one billion people globally.

The Flinders Health and Medical Research Institute (FHMRI) Sleep Health research is one of the first large-scale extended studies of OSA severity, made possible by using a novel under-mattress sleep sensor technology that allows for multi-night, in-home, non-invasive monitoring of sleep data from 12,287 adults over approximately 180 nights with about 30 repeat blood pressure tests.

The sensor technology measured both OSA severity and night-to-night variation in OSA severity, leading researchers to make comparisons with current single night sleep assessments and possible misdirected management and care.

"These findings reinforce recent evidence indicating there is considerable night-to-night variation in OSA severity for many people, which raises concerns clinically regarding OSA misdiagnosis," says

senior Flinders University sleep researcher and FHMRI Sleep Health director, Professor Danny Eckert.

"Furthermore, our study suggests that high night-to-night variability in OSA severity may also be an important contributor to cardiovascular disease—one of the world's leading causes of death."

"Therefore, there is considerable potential to incorporate new simplified monitoring approaches to aid current single-night diagnostics, which are labor intensive and costly."

More information: Bastien Lechat et al, High night-to-night variability in sleep apnea severity is associated with uncontrolled hypertension, *npj Digital Medicine* (2023). [DOI: 10.1038/s41746-023-00801-2](https://doi.org/10.1038/s41746-023-00801-2)

Provided by Flinders University

Citation: Nightly variation in sleep apnea sharply increases cardio risks, shows study (2023, April 26) retrieved 14 May 2024 from

<https://medicalxpress.com/news/2023-04-nightly-variation-apnea-sharply-cardio.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--