


Study shows pediatric health professionals had accentuated psychological disorders during the COVID-19 pandemic

April 21 2023



Study Flowchart. Credit: *Pediatric Critical Care Medicine* (2022). DOI: 10.1097/PCC.00000000000003119

Concern about the mental health of hospital professionals has been increasing in recent years, and when, in early 2020, the COVID-19

pandemic was declared and rapidly spread, there was a large increase in the overload of workers in Intensive Care Units (ICUs).

Given this, researchers from the D'Or Institute for Research and Education (IDOR) and other national institutions assessed the prevalence and extent of psychological disorders such as burnout, anxiety, depression, and [post-traumatic stress disorder](#) (PTSD) in pediatric ICUs (PICUs) [health professionals](#). The study was published in the scientific journal *Pediatric Critical Care Medicine* and selected as the Feature Article and Editor's choice of the month, receiving a specific editorial.

In the editorial published about the study, Dr. Gillian Colville starts by asking, "How was work today?" and points out, "It is an innocent question, but not an easy one to answer when you work in intensive care. Aside from obvious issues relating to confidentiality, so much of what we witness working in these settings is unspeakable, shocking, and sad. And when the unit is one that treats children, it feels socially unacceptable, outside the hospital, to even hint at what we are dealing with daily."

The COVID-19 pandemic imposed an exhaustive and prolonged workload on health professionals, especially those in the ICU, due to the clinical condition of the patients and, as highlighted by Dr. Fernanda Lima-Setta, the first author of the study, "Pediatric ICU workers also faced the biggest challenges, even though children were not the age group most seriously affected in the pandemic. There were many changes in clinical protocols, and the team had to deal with the multisystem inflammatory syndrome associated with COVID-19, which had a [fatality rate](#) of 6.85%."

Furthermore, children remain a sensitive group in many countries due to the delay in the progress of pediatric vaccination.

This research included 29 PICUs (13 public and 16 private) in different Brazilian regions between July and October 2020. Participants received an [online survey](#) that encompassed demographic variables, occupational characteristics, and self-reported mental and emotional well-being questionnaires. Among the professionals eligible for the study, 1,084 answered the questionnaire, which included physicians, nurses, physiotherapists, and nursing technicians.

The respondents were mostly young, female, and one-third were physicians. About 20% of the participants reported a previous mental health disorder diagnosis confirmed by a specialist, and almost 80% declared themselves satisfied or very satisfied with their current job. On the questionnaire, burnout scores concerning emotional exhaustion and personal achievement were high.

The prevalence of anxiety (33%), depression (19%), and PTSD (13%) were also found. In addition, the researchers also noted that those with prior mental health issues scored the highest, and that younger staff members reported more burnout.

It is worth noting that professionals working in the public health system were more exposed to the risk of burnout and PTSD. In Brazil, public health institutions are responsible for most of the population's care, and these have a chronic scarcity of resources and labor, culminating in a much greater workload and stress for the PICU team.

In the professional categories, physicians had worse burnout scores, although no differences in prevalence were found between subgroups. Complex decision-making when dealing with limited resources in the face of the extremely high demands imposed by the pandemic may have influenced this data.

The authors report that there are few studies related to pediatrics to

make a comparison possible. However, burnout rates were higher than those obtained in another study conducted by a group of IDOR researchers before the pandemic, involving adult ICU professionals. A limitation point of this research is the fact that it is a cross-sectional study, and it is not possible to assess the cause-effect relationship.

Brazilian PICU workers appear to be at greater risk of experiencing symptoms of mental disorders during the first peak of COVID-19 in 2020 compared to the pre-pandemic period. Despite high levels of personal achievement combined with general satisfaction with the current job, organizational support can help in challenging situations of psychological distress and traumatic stress.

In the editorial, Dr. Colville concluded that "We need to be able to answer honestly 'How was work today?' and to acknowledge that even in the darkest of times, it is possible to find meaning in what we do."

More information: Fernanda Lima-Setta et al, Mental Health and Emotional Disorders During the COVID-19 Pandemics: Prevalence and Extent in PICU Staff*, *Pediatric Critical Care Medicine* (2022). [DOI: 10.1097/PCC.0000000000003119](https://doi.org/10.1097/PCC.0000000000003119)

Provided by D'Or Institute for Research and Education

Citation: Study shows pediatric health professionals had accentuated psychological disorders during the COVID-19 pandemic (2023, April 21) retrieved 5 May 2024 from <https://medicalxpress.com/news/2023-04-pediatric-health-professionals-accentuated-psychological.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
