


# A peek into the interplay between vision and decision making in the brain

April 19 2023



Choice is distributed, near orthogonal to other components and with a ventral-stream dominance. **a** Angles between state axes averaged across animals. Choice axes (pre- and post-movement) were orthogonal to all other axes (smallest angle  $84 \pm 7^\circ$ ). Attention and wheel had the smallest angular separation ( $44 \pm 3^\circ$ ), followed by wheel and saccades ( $56 \pm 4^\circ$ ). **b** Hierarchical clustering from the angle distances in **a**. Attention and wheel movements were most similar. Choice pre- and post-movement onset clustered together, whereas stimulus and saccades had unique profiles. **c** Angle between choice and movement state axes averaged across animals aligned to movement onset (line for average across animals and shaded area for its 95% CI,  $n = 7$  animals); shaded gray band is the expected angular distance range for statistically independent axes; observed angles never

significantly deviated from the statistically independent condition. d Spatial-Distribution index (SDI) for each state axis. Choice had the largest SDI ( $30 \pm 4\%$ ); dots are different animals; middle lines and shaded areas are means and their 95% CI ( $n = 7$  animals). e We computed five  $d'$  values, each derived by restricting locaNMF components to one of the five area groups (insets), thus defining a 5-D space for  $d'$  components. The five broad area groups consisted of the dorsal stream (PM and AM), ventral stream (L), posterior parietal (A, AL, and RL), somatosensory (SS<sub>t</sub> and SS<sub>b</sub>) and retrosplenial (RS) regions. Each plot shows a 2-D projection of the five broad area groups, where each dot corresponds to the  $d'$  values for a given animal and the large dot with errorbars to the average across animals and s.e. ( $n = 7$  animals). Credit: *Nature Communications* (2023). DOI: 10.1038/s41467-023-35824-6

An all-RIKEN team of neuroscientists has mapped out neurons in the mouse visual system that are selectively activated during decision-making, and may directly influence this process. To achieve this, they combined a cleverly-crafted behavioral task with sophisticated imaging of neurons.

The choices we make involve diverse sectors of the [brain](#), including those responsible for cognition, movement and other activities. This complexity has made it challenging for neuroscientists to tease out which [neural circuits](#) directly coordinate decision making.

Many [important decisions](#) start with responding to what we see in front of us. But the visual centers of the mammalian brain rely on much more information than is supplied by the eyes alone. "Most of the inputs come from within the brain," says Andrea Benucci of the RIKEN Center for Brain Science. "And they carry a great diversity of non-visual signals."

This led Benucci to hypothesize that these other sources of data may represent choice-guiding neuronal inputs. However, confirming this

would require designing an experiment that minimized the confounding effects of other brain functions that intersect with the [decision-making](#) process.

To tackle this challenge, Benucci and his team used a behavioral task for mice that minimized the involvement of memory, [body movements](#) and responses to novel stimuli. This entailed training animals to earn a water reward by aligning the orientation of a projected image to match that of a target pattern. Once the animals had mastered this task, the researchers used a fluorescent "reporter" protein to monitor the firing of neurons in different regions of the brain as animals pursued their reward.

Their imaging data revealed subsets of neurons that appeared to be directly involved in choice. These were mostly located in regions of the brain within the ventral visual stream, a system of neurons primarily involved in identifying features in a visual scene. The strength of these signals depended both on the difficulty of the task and on the extent to which the animal's attention was focused on the orientation task.

Benucci and colleagues validated the activity patterns they had observed with a neural-network-based [computational model](#), which they trained using the actual data from animals. The work is published in the journal *Nature Communications*.

"In this study, we not only uncovered signatures of choice, but we also found well-structured representational dynamics of choice signals that reflected the context-dependent nature of the decision-making process," says Benucci. "The model confirmed that the recorded signals indeed reflect network computations associated with solving this [specific task](#) using choice strategies matching those of the mice."

Intriguingly, the choice signals identified here were produced by a relatively small number of sparsely dispersed neurons within the ventral

stream. Benucci's team is now exploring the extent to which these "loners" influence vision-based decision making.

**More information:** Javier G. Orlandi et al, Distributed context-dependent choice information in mouse posterior cortex, *Nature Communications* (2023). [DOI: 10.1038/s41467-023-35824-6](https://doi.org/10.1038/s41467-023-35824-6)

Provided by RIKEN

Citation: A peek into the interplay between vision and decision making in the brain (2023, April 19) retrieved 11 May 2024 from <https://medicalxpress.com/news/2023-04-peek-interplay-vision-decision-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.