

Severe COVID-19 linked with 16-fold risk of life-threatening heart rhythm within 6 months

April 18 2023



Credit: Unsplash/CC0 Public Domain

Patients with severe COVID-19 requiring mechanical ventilation are 16 times more likely to develop ventricular tachycardia within six months compared to their peers without severe infection, according to research presented at <u>EHRA 2023</u>, a scientific congress of the European Society



of Cardiology (ESC). Risks of other heart rhythm disorders were also elevated.

"The actual likelihood of developing ventricular tachycardia or other arrhythmias after severe COVID-19 is low for the individual patient, but much higher than in those without severe infection," said study author Dr. Marcus Stahlberg of the Karolinska Institute, Stockholm, Sweden.

The objective of this study was to investigate the long-term risk of arrhythmias after discharge from an <u>intensive care unit</u> (ICU) for COVID-19 requiring mechanical ventilation. The researchers used the Swedish ICU register to identify all COVID-19 patients treated with mechanical ventilation and discharged alive from an ICU between March 2020 and June 2021.

Each patient was matched by age, sex and district of residence with up to 10 people in the general population. Multiple compulsory national registries were used to record new diagnoses of arrhythmias after discharge from the ICU.

The primary outcome was hospitalization with ventricular tachycardia, atrial fibrillation, other tachyarrhythmias, or bradycardia/pacemaker implantation. Ventricular tachycardia is a potentially life-threatening heart rhythm disturbance that occurs when the ventricle beats too fast to pump properly causing the body to receive insufficient oxygenated blood. Atrial fibrillation is an irregular and fast heartbeat that causes shortness of breath and raises the risk of stroke.

"Other tachyarrhythmias" refers to a fast heartbeat not caused by atrial fibrillation. "Bradycardia or pacemaker implantation" is a combined endpoint of slow heart rate or need for a pacemaker due to slow heart rate.



The researchers analyzed the risk of developing each <u>arrhythmia</u> in severe COVID-19 patients, compared to those without severe COVID-19. The analyses were adjusted for factors linked with the likelihood of heart rhythm disorders including age, sex, <u>high blood</u> <u>pressure</u>, diabetes, high blood lipids, <u>chronic kidney disease</u> and <u>socioeconomic status</u> (education level, marital status and income).

The study included 3,023 patients with severe COVID-19 who received mechanical ventilation at a Swedish ICU and 28,463 individuals from the general population who had not been in an ICU with COVID-19 requiring mechanical ventilation (control group). The average age of participants was 62 years and 30% were women. "Higher age and male sex are two important risk factors for getting severely sick with COVID-19 and this was reflected in our study participants," said Dr. Stahlberg.

The average follow up was nine months. In patients who had severe COVID-19, the incidence rates per 1,000 person-years of ventricular tachycardia, atrial fibrillation, other tachyarrhythmias, and bradycardia/pacemaker implantation were 15.4, 78.4, 99.3 and 8.5, respectively. Corresponding incidence rates in the control group were 0.9, 6.0, 6.7 and 0.9, respectively.

Compared to the control group, patients who had severe COVID-19 requiring mechanical ventilation had a 16-fold risk of ventricular tachycardia, 13-fold risk of <u>atrial fibrillation</u>, 14-fold risk of other tachyarrhythmias, and 9-fold risk of bradycardia/pacemaker implantation.

Dr. Stahlberg said, "COVID-19 patients who need <u>mechanical</u> <u>ventilation</u> often have other conditions and adding a heart rhythm disorder may lead to worsened health. These patients should seek medical attention if they develop palpitations or irregular heartbeats



after hospital discharge so they can be evaluated for possible arrhythmias."

He concluded, "An increased risk of arrhythmias following COVID-19 has also been reported previously in the bulk of COVID-19 patients not requiring ICU treatment. Together with our new data and taking into the account that we globally have 650 million reported COVID-19 cases, hospital systems should prepare for an increase in patients requiring management for new onset arrhythmias."

Provided by European Society of Cardiology

Citation: Severe COVID-19 linked with 16-fold risk of life-threatening heart rhythm within 6 months (2023, April 18) retrieved 4 May 2024 from https://medicalxpress.com/news/2023-04-severe-covid-linked-fold-life-threatening.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.