


Surgery is the most effective treatment for metabolic liver disease, says new study

April 20 2023



Credit: CC0 Public Domain

Metabolic (bariatric) surgery is more effective than medications and lifestyle interventions for the treatment of advanced non-alcoholic fatty liver disease.

A new paper, published today in *The Lancet* by King's College London and the Catholic University of Rome, is the first to compare three active treatments of [non-alcoholic steatohepatitis](#) (NASH) and to specifically investigate the effectiveness of [metabolic surgery](#) (weight loss surgery) in a randomized clinical trial.

Non-alcoholic fatty liver disease (NAFLD) is the most common cause of chronic liver disease, globally affecting 55% of people with type 2 diabetes and 75% of those with obesity. Non-alcoholic steatohepatitis (NASH) is the progressive form of the disease, and is characterized by liver cell injury and inflammation, which induce liver fibrosis (scarring of the tissue). Left untreated, it can lead to [liver failure](#) and [liver cancer](#), and is one of the leading causes of liver transplant in the western world.

NASH also increases risk of cardiovascular disease and overall mortality in patients with obesity or type 2 diabetes. NAFLD/NASH currently affects 12% of adults in the UK; in the U.S. it is estimated that the condition will affect 27 million people by 2030.

The multi-center randomized trial was conducted in Italy. The trial compared the efficacy of bariatric and metabolic surgery versus lifestyle modifications and best current medical care in 288 patients.

Pre- and post-operative liver biopsies showed that surgery was more effective in inducing complete reversal of inflammation and cell damage

in the liver—the core characteristics of NASH—without worsening of liver fibrosis after 1 year from surgery. The probability of achieving reversal of NASH was 3–5 times higher with metabolic surgery than with medical care. Surgery was also more effective at achieving improvement of at least one stage of [liver fibrosis](#), another pre-specified endpoint of the trial. The two surgical procedures appeared to equally improve NASH.

Researchers say the ability of surgery to control and even improve fibrosis associated with NASH is of special clinical relevance, as fibrosis is the main predictor of liver complications and poor cardiovascular outcomes and death in patients with NASH.

Professor Geltrude Mingrone, first author of the report, Professor of Medicine at the Catholic University of Rome and a Professor of Diabetes and Nutrition at King's College London, said, "The results of our study support the use of metabolic surgery as a treatment of NASH, a condition for long considered orphan of effective therapies."

Professor Francesco Rubino, a senior co-investigator and Chair of Bariatric and Metabolic Surgery at King's College London and consultant surgeon at King's College Hospital, said, "The presence of NASH predicts a significant risk of complications and mortality in people with severe obesity and type 2 diabetes. The results of this study provide a compelling case for prioritization of metabolic surgery in this patient population."

Compared to conventional medical treatment, [surgery](#) also resulted in better overall health benefits. While both surgical procedures seemed to equally improve NASH, the gastric bypass group was more effective than [sleeve gastrectomy](#) at improving type 2 diabetes and reducing other cardiovascular risk factors associated with NAFLD/NASH.

More information: Bariatric–metabolic surgery versus lifestyle intervention plus best medical care in non-alcoholic steatohepatitis (BRAVES): a multicentre, open-label, randomised trial, *The Lancet* (2023).

Provided by King's College London

Citation: Surgery is the most effective treatment for metabolic liver disease, says new study (2023, April 20) retrieved 24 May 2024 from <https://medicalxpress.com/news/2023-04-surgery-effective-treatment-metabolic-liver.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--