


Amino acid PET successfully differentiates recurrent brain metastases, reducing invasive procedures and overtreatment

May 12 2023



Methods

- 12 studies with tracers
- FET, MET, FDOPA
- 547 lesions in 397 patients

Results

- Pooled sensitivity: 82%
- Pooled specificity: 84%
- Odds ratio: 17.1

Conclusions

Good accuracy of amino acid PET in the differential diagnosis of recurrent brain metastases

Does a new or increasing contrast enhancement on T1 MRI represent a consequence of treatment or progressive/recurrent tumor? This common clinical dilemma can be resolved by amino acid PET with good diagnostic accuracy.

Credit: Society of Nuclear Medicine

A newly published meta-analysis indicates that amino acid PET can accurately differentiate recurrent or progressive brain metastases from

treatment-related changes. A specificity of 84% suggests that it may reduce the number of invasive procedures and overtreatment in patients who in fact experience treatment-related changes. This research was published in the May issue of the *Journal of Nuclear Medicine*.

Brain metastases occur in 20% to 40% of all [cancer patients](#) and are most likely to occur in those with lung, breast and renal cancer, melanoma, and cancers of the gastrointestinal tract. Management of [patients](#) with [brain metastases](#) usually includes surgery, radiation and chemotherapy. Some patients develop treatment-related changes such as radiation necrosis or pseudoprogression.

"A differentiation between recurrent or progressive brain metastases and treatment-related changes is challenging," said Igor Yakushev, senior physician in the department of nuclear medicine at Technical University of Munich in Germany. "As the management of patients with recurrent or progressive brain metastases and treatment-related changes is fairly different, accurate and early differential diagnosis is essential."

The [meta-analysis](#) included 12 studies with amino acid PET radiotracers. The studies included a total of 397 patients with 547 lesions. Overall, 269 lesions (49%) were found to be recurrent or progressive brain metastases. Using a histologic examination and/or radiological and clinical follow-up as reference, pooled sensitivity and specificity of amino acid PET were found to be 82% and 84% respectively.

"This study provides IIa class evidence on diagnostic utility of amino acid PET in the differential diagnosis of recurrent or progressive brain metastases," stated Yakushev. "These findings are in line with an increasing role of molecular imaging in the management of patients with [brain tumors](#), yet the results also point to potential for further improvement of diagnostic accuracy."

More information: Timo Schlürmann et al, Utility of Amino Acid PET in the Differential Diagnosis of Recurrent Brain Metastases and Treatment-Related Changes: A Meta-analysis, *Journal of Nuclear Medicine* (2022). [DOI: 10.2967/jnumed.122.264803](https://doi.org/10.2967/jnumed.122.264803)

Provided by Society of Nuclear Medicine

Citation: Amino acid PET successfully differentiates recurrent brain metastases, reducing invasive procedures and overtreatment (2023, May 12) retrieved 5 May 2024 from <https://medicalxpress.com/news/2023-05-amino-acid-pet-successfully-differentiates.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--