

# Combination therapy outperforms chemotherapy in clinical trial for patients with chronic lymphocytic leukemia

May 10 2023

---



Credit: Unsplash/CC0 Public Domain

Chemotherapy has been the most effective treatment to date for young and fit patients with chronic lymphocytic leukemia (CLL). The disease

is the most common form of leukemia in the Western world, causing a substantial health burden on patients and society. It is currently incurable. Research led by researchers from Cologne University, Amsterdam UMC and Copenhagen University now shows that a chemo-free drug-combination therapy is more effective and produces fewer side effects. The results of this study are published today in the *New England Journal of Medicine*.

Research at Amsterdam UMC in recent years has contributed greatly to a better understanding of the biology of the disease. This study, the largest ever conducted in CLL, with a younger and fitter patient group builds upon these findings. CLL originates in B cells, a type of white blood cell that produces antibodies against bacteria and viruses. Therefore, one of the most common complications of this disease is infections, which are often severe.

Previously it was believed that these CLLS were very difficult to kill. However, this was found not to be the case, granting researchers access to a raft of new treatment options. Arnon Kater, professor of hematology at Amsterdam UMC and chair of the HOVON CLL study group, explains that "just like a Formula 1 car on the track, we saw that the CLL cells in the blood are very vulnerable. The protein Bcl-2 protects cells from dying in the blood. This protection gives the cells time to go for a pit stop, just like a Formula 1 car. The pit stop is the lymph node where the cancer cells can recharge and then divide."

The researchers learned from previous studies that the drug [venetoclax](#) effectively inhibits the Bcl-2 protein. Cells in the blood do still die, but crucially, not during the pit stop in the lymph node.

"By combining venetoclax with drugs that can do their job within the pit lane, we thought that combinations were possible that were debilitating and would be more effective than chemotherapy," says Kater.

## Largest study to date

The phase 3 randomized trial, also known as the GAIA/CLL13 trial, was conducted with a total of 920 fit patients with CLL in 159 hospitals in nine European countries and Israel. About a quarter of the patients came from the Netherlands. The patients were placed in four groups and received either chemotherapy and an antibody (standard treatment), or the drug venetoclax in combination with an antibody (either rituximab or [obinutuzumab](#)). The fourth group received venetoclax, obinutuzumab and the specific kinase inhibitor ibrutinib.

Importantly, in all four groups, the treatment was temporary, in contrast to earlier studies that continued either venetoclax or ibrutinib, which prolongs exposure to side effects, dramatically increases the costs and inevitably results in development of resistance.

All groups with venetoclax had fewer side effects, and that the combinations of venetoclax with obinutuzumab were more beneficial than the [standard treatment](#) with chemotherapy. Treatment was specifically discontinued in patients with an aggressive form of the disease (so-called IGHV unmutated variant, which resolves in 50% of patients).

On average, after 3 years in the more aggressive disease group, in 35% of those who received chemotherapy, the disease had returned. For those who received the experimental combination therapy, this was only 18%. Making the combination therapy twice as effective for this group.

## Benefits of combination therapy

Until now, chemo-immunotherapy has been the most effective treatment for patients with treatment continuing indefinitely. With drug-

combination therapy it is now possible to stop treatment after a year.

"This study shows that with clever temporary and safe combinations, you can allow patients to be treatment free in the long-term, with a much lower chance of developing resistance. And we also think that it is even possible to stop the combination therapy earlier than after a year. We now want to investigate this," says Kater. This not only reduces side effects, but also health care costs.

**More information:** First-Line Venetoclax Combinations in Chronic Lymphocytic Leukemia, *New England Journal of Medicine* (2023). [DOI: 10.1056/NEJMoa2213093](https://doi.org/10.1056/NEJMoa2213093)

Provided by University of Amsterdam

Citation: Combination therapy outperforms chemotherapy in clinical trial for patients with chronic lymphocytic leukemia (2023, May 10) retrieved 30 June 2024 from <https://medicalxpress.com/news/2023-05-combination-therapy-outperforms-chemotherapy-clinical.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.