


## Risk factors for Parkinson's disease: Study proves the protective function of the RIT2 gene

May 25 2023



In vivo aSyn overexpression increases endogenous LRRK2 activity, which is prevented by Rit2 co-expression. **a** PLA analysis of AAV-GFP, AAV-A53T-aSyn and AAV-A53T-aSyn + AAV-Rit2 injected mice in TH-positive neurons in the SNc. **b** Quantification of PLA counts in TH-positive neurons shows a significant increase of endogenous LRRK2 kinase activity with AAV-A53T-aSyn injection. The increase is completely prevented by co-injection of AAV-



Rit2 with AAV-A53T-aSyn (AAV-GFP = 6 animals, AAV-A53T-aSyn = 6 animals, AAV-A53T-aSyn + AAV-Rit2 = 5 animals). Data represented as mean  $\pm$  SEM. \*p

Citation: Risk factors for Parkinson's disease: Study proves the protective function of the RIT2 gene (2023, May 25) retrieved 27 April 2024 from <a href="https://medicalxpress.com/news/2023-05-factors-parkinson-disease-function-rit2.html">https://medicalxpress.com/news/2023-05-factors-parkinson-disease-function-rit2.html</a>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.