

Inhibiting the biological crosstalk of autophagy and mitochondrial function underlying pancreatic ductal adenocarcinoma

May 5 2023, by Thamarasee Jeewandara



Autophagy inhibition leads to a drop in the LIP in PDAC. (A) Autophagy was inhibited genetically or pharmacologically in PDAC cells, and the relative LIP was determined after cotreatment with FAC in 8988T cells. (B) Chloroquine (CQ)–mediated drop in LIP was rescued by FAC in the indicated PDAC cell lines. (C) OCR rescue after FAC cotreatment in CQ-treated or ATG5-knockdown PDAC cells. Heatmaps showing clonogenic assays (D) in



PDAC cells and relative proliferation in a panel of PDAC cell lines after treatment with FAC in autophagy-inhibited cells. Data are means ± SD, and P values were quantified using two-way analysis of variance (ANOVA) with Sidak's multiple comparison's test (for A, B, D, and E) and one-way ANOVA with Tukey's post hoc test (for C). **P Science Advances (2023). DOI: 10.1126/sciadv.adf9284

Pancreatic ductal adenocarcinoma (PDAC) cells maintain a high level of autophagy or degradation, allowing them to thrive in severely limiting microenvironments. However, the process via which <u>autophagy</u> promotes pancreatic cancer cell growth and survival have yet to be understood.

In a new report published in *Science Advances*, Subhadip Mukhopadhyay and a research team in <u>radiation oncology</u> and genome stability at the NYU and Harvard, showed how the inhibition of autophagy altered the mitochondrial function in PDAC patients. They experimentally attenuated the <u>succinate dehydrogenase complex iron sulfur subunit</u> in mitochondria to limit the availability of the labile iron pool.

The cancer cell lines used autophagy to sustain iron homeostasis in contrast to other <u>tumor types</u> that relied on <u>macropinocytosis</u>. The biologists noted how cancer-associated fibroblasts provided bioavailable iron to pancreatic <u>cancer cells</u> to resist the ablation of autophagy. Mukhopadhyay and colleagues administered a low-iron diet in a <u>mouse</u> <u>model</u> to hinder this crosstalk in the form of autophagy inhibition therapy. The outcomes emphasized a critical link between autophagy, iron metabolism and mitochondrial function to drive the progression of pancreatic ductal adenocarcinoma.

Iron homeostasis in pancreatic ductal



adenocarcinoma (PDAC)

Pancreatic ductal adenocarcinoma represents more than <u>95% of</u> pancreatic cancers that are highly resistant to therapy with a low 5-year survival rate, where the number of cancer-related deaths accounted to the second highest rate in the United States. These tumors are defined by a harsh, hypoperfused and hypoxic microenvironment with altered availability of nutrients. The cancer cells can <u>survive hostile</u> <u>environments</u> by reprogramming their metabolic needs to rely on nutrient scavenging mechanisms such as macroautophagy and macropinocytosis to survive and grow. Previous work had highlighted the role of <u>ferritinophagy</u>—a selective form of autophagy—and its role in <u>iron homeostasis</u> maintenance.





Autophagy inhibition abrogates SDHB level in PDAC. Pharmacological (A) or genetic (B) inhibition of autophagy in PDAC cells was cotreated with FAC for 24 hours followed by immunoblots for the indicated proteins. SDHB is pointed out using a triangle. Similar to (A), other PDAC cell lines were used for analyzing SDHB level in (C) followed by total ion counts of succinate in PDAC cells using liquid chromatography–mass spectrometry (n = 3 technical replicates) after autophagy inhibition (D). Data are means ± SD, and P values were quantified using unpaired Student's two tailed t test. *P Science Advances (2023). DOI: 10.1126/sciadv.adf9284



In this work, Mukhopadhyay and the team showed how the inhibition of autophagy reduced the assembly of iron-sulfur clusters to stabilize a variety of proteins, including those involved in the <u>electron transfer</u> chain, such as <u>succinate dehydrogenase complex iron sulfur subunit B</u> (SDHB). The reduction of this molecule in pancreatic cancer cells during autophagy led to a decrease in mitochondria respiration and reorganization of the cristae microarchitecture. The supplementation of iron or the expression of ectopic SDHB revived the growth and improved the mitochondrial defects.

Limiting the bioavailable iron in pancreatic ductile cancer cells to inhibit autophagy

Previously, Mukhopadhyay and colleagues showed how autophagy inhibition in pancreatic <u>cancer cell lines</u> led to a decrease in the mitochondrial oxygen consumption rate. However, they could not determine the mechanism of this impairment. Prior research had shown amino acid pools to remain unaltered upon autophagy inhibition, <u>except</u> <u>for a decrease in cysteine</u> levels.

However, supplementing cysteine to pancreatic cancer cells did not alter the defective rate of oxygen consumption, indicating a pathway that directly affected the mitochondria. While iron-sulfur cluster proteins drove mitochondrial function, a drop in the labile iron pool after autophagy inhibition decreased mitochondrial respiration, which the team restored via <u>ferric ammonium citrate supplementation</u>. The inhibition of autophagy further led to decreased levels of the succinate dehydrogenase complex iron sulfur subunit B (SDHB) protein, highlighting a functional loss of the mitochondrial protein.





Loss of autophagy impairs mitochondrial microarchitecture in PDAC. (A) Representative TEM images of PDAC cells treated with PBS (Control) or CQ followed by determining their (B) mitochondrial number per cell (Control, n = 14; CQ, n = 12 cells). (C) A magnified section of a PDAC cell showing mitochondrial ultrastructure after CQ treatment followed by quantification of (D) crista number per mitochondria (n = 17 unique crista for both Control and CQ), (E) crista lumen width (Control, n = 18; CQ, n = 13 unique crista), and (F) crista length (n = 19 unique crista for both Control and CQ). (G) Representative TEM images of ATG5-knockdown PDAC cells with or without FAC treatment were quantified for (H) crista length and (I) crista lumen width (siControl, n = 13; siATG5, n = 9; siATG5 + FAC, n = 11 unique crista). Black boxes were



digitally zoomed in to highlight the morphology of typical dysfunctional mitochondria. Random TEM image of mitochondria was blindly acquired and quantified. Data are means ± SD, and P values were quantified using one-way ANOVA with Tukey's post hoc test. **P Science Advances (2023). DOI: 10.1126/sciadv.adf9284

Defects in the mitochondrial architecture are related to autophagy inhibition

The research team used <u>transmission electron microscopy</u> to reveal the mitochondrial architecture of autophagy-deficient pancreatic cancer cells; enabled by administering chloroquine. They noted a significant reduction in mitochondrial cristae lumen with large vacant spaces post-treatment to indicate functional defects.

The team showed how the ectopic expression of the SDHB protein could restore the cristae microarchitecture. In fact, overexpression of SDHB rescued the proliferation and oxygen consumption rate after autophagy inhibition. They studied the mechanism via which these components played a significant role during the biogenesis of iron and sulfur proteins, after the loss of autophagy in pancreatic cancer cells.

Investigating the crosstalk between cancer-associated cell co-cultures with drug moieties

While autophagy can regulate the availability of iron in pancreatic cancer cells, the patient response to <u>hydroxychloroquine monotherapy</u> was not clinically sufficient to attenuate the disease. To examine the impact of the tumor microenvironment to this resistance, the researchers studied the effects of compensating the labile iron pool by co-culturing pancreatic cancer cells with cancer-associated fibroblasts to recreate a



microphysiological tumor environment.

The researchers studied the crosstalk between cells to understand cellular regulation of the labile iron pool and examined the expression of key iron transporters. They noted an increase in the exporter ferroprotein in fibroblasts cultured alongside autophagy-inhibited pancreatic cancer cells. When they administered <u>Tocilizumab</u>, they restored the antiproliferative effects of autophagy inhibition on the cancer cell line in the co-culture system.



Ectopic SDHB rescues the mitochondrial dysfunction upon autophagy inhibition in PDAC. PDAC cells expressing SDHB and control vector were treated with CQ followed by immunoblotting for the indicated proteins (A) and analyzed for (B) OCR (combined data from n = 2 experiments). (C) Representative TEM images of cells are shown in situation similar to (A) followed by mitochondrial



crista length (D; n = 18 unique cristae for all groups) and lumen width (E; n = 18 unique cristae for all groups) determination. Red arrows in (C) indicate the typical dysfunctional mitochondria. ATG5 was suppressed in SDHB overexpression and control PDAC cells for immunoblotting of indicated proteins (F). Under conditions similar to (F), cell proliferation (G; combined data of n = 5 experiment) and OCR was determined (H; combined data of n = 2 experiments). Data are means \pm SD, and P values were quantified using one-way ANOVA with Tukey's post hoc test. *P Science Advances (2023). DOI: 10.1126/sciadv.adf9284

Treating pancreatic ductal adenocarcinoma (PDAC)

Mukhopadhyay and colleagues explored the methods to treat the cancer and explored if autophagy inhibition could facilitate iron restriction to inhibit pancreatic tumor growth. They decreased the bioavailable iron to impair the crosstalk between <u>pancreatic duct adenocarcinoma cells</u> and <u>cancer-associated fibroblasts</u>. They accomplished this by systematically reducing iron levels in mice by exposing them to a low-iron diet for two weeks before transplanting cancer cells with a <u>doxycycline-inducible</u> <u>dominant vector</u> and a control vector in mouse hosts. While the iron level in the diet did not affect the level of hemoglobin, autophagy inhibition significantly decreased tumor growth. The low-iron diet further reduced tumor growth to suggest that iron deprivation sensitized tumor cells to undergo autophagy inhibition.

The combination of an iron-limiting diet and autophagy decreased the bioavailable ferrous iron concentration in tumors, while decreasing the succinate dehydrogenase complex iron sulfur subunit B expression. When the scientists administered chloroquine alone, it failed to reduce tumor growth in mice, although combined chloroquine and an iron-restricted diet significantly reduced tumor growth. The combined data represented a low-iron diet that worked together with autophagy



inhibition to critically disrupt iron homeostasis in pancreatic <u>ductal</u> <u>adenocarcinoma</u> tumors.



Combination of iron restricted diet and autophagy inhibition provides a therapeutic benefit in PDAC. (A) Murine PDAC cells (HY19636) bearing intact autophagy (mSt) or inhibited autophagy (4B) were orthotopically implanted in syngeneic mice fed with doxycycline (625 mg/kg) diet containing control (220 ppm) or low iron (4 to 6 ppm). Mice were euthanized after 24 days of implantation, and tumor images are shown (A; n = 10, 9, 10, and 10 from top to bottom) along with tumor weights (B). (C) The ferrous concentration of different tumors indicated in (A) was analyzed from three randomly selected tumors of each group. (D) SDHB expression levels were determined by performing immunoblot analysis on these tumors, and their relative quantification was performed (E). B6 mice bearing syngeneic PDAC cells were treated with CQ or PBS, fed with control or low-iron diet, and were euthanized 24 days after implantation. Tumor images are shown (F; n = 12, 10, 12, and 12



from top to bottom) along with tumor weights (G). (H) Ferrous concentration in tumors of the indicated groups was determined (n = 3). (I) SDHB expression level was assessed by immunoblotting on these tumors, and their relative quantification was performed (J) (n = 3). Data are means \pm SD, and P values were quantified using one-way ANOVA with Tukey's post hoc test. *P Science Advances (2023). DOI: 10.1126/sciadv.adf9284

Outlook

In this way, Subhadip Mukhopadhyay and colleagues identified how pancreatic duct adenocarcinoma tumors relied on <u>autophagy</u> to grow and evolve with time during disease progression. Autophagy inhibition decreased the labile iron pool and the synthesis of iron-sulfur clusters. They supplemented the cultures with iron and promoted the ectopic expression of both iron-sulfur clusters and succinate dehydrogenase complex iron sulfur subunit B (SDHB) protein to restore mitochondria function and cell growth. Autophagy was essential to maintain iron homeostasis, and support <u>mitochondrial function</u>, required for both growth and proliferation, in pancreatic cancer cells.

The researchers dissected multiple metabolic cross-talks occurring between the tumor and stroma, and explored the disruption of iron metabolism in the pancreatic duct adenocarcinoma cells and its impact on the complex tumor microenvironment. These mechanistic findings will assist in the development of new design approaches to target the process of <u>iron</u> metabolism in pancreatic duct adenocarcinoma cells.

More information: Subhadip Mukhopadhyay et al, Autophagy supports mitochondrial metabolism through the regulation of iron homeostasis in pancreatic cancer, *Science Advances* (2023). DOI: 10.1126/sciadv.adf9284



Cristovão M. Sousa et al, Pancreatic stellate cells support tumour metabolism through autophagic alanine secretion, *Nature* (2016). <u>DOI:</u> <u>10.1038/nature19084</u>

© 2023 Science X Network

Citation: Inhibiting the biological crosstalk of autophagy and mitochondrial function underlying pancreatic ductal adenocarcinoma (2023, May 5) retrieved 4 May 2024 from https://medicalxpress.com/news/2023-05-inhibiting-biological-crosstalk-autophagy-mitochondrial.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.