

Genetics and concussion—why a minor knock can be devastating for some people

June 20 2023, by Lyn Griffiths and Annette Greenhow



Credit: AI-generated image ([disclaimer](#))

Concussion and head trauma is a real and serious risk for many Australians. While most people suffer acute and relatively short-lived effects, such as dizziness and headache, in some cases symptoms persist for weeks, months or years. It can result in long-term and debilitating neurological impairment.

Concussion in sport—from the junior to the elite level—is being prioritized as a public health concern in Australia. A [Senate inquiry](#) into concussions and repeated head trauma in contact sport is due to report in August. Of note in the hearings has been the AFL's [acknowledgement](#) of an association between head trauma and [chronic traumatic encephalopathy](#), a neurodegenerative disease [found](#) in several deceased players.

The [latest data](#) show [concussion](#) can happen in nearly every sport, not just contact sports, with almost 3,100 hospitalizations for concussion caused by sports in 2020–21.

But not everyone responds the same way to concussion. At present, there are [few reliable indicators](#) of who will suffer specific or long-term effects. We do know the number and severity of [symptoms](#) and [multiple concussions](#) are important. And we are developing understanding of how a person's genes play a role.

Traumatic brain injury

Concussion is a form of traumatic brain injury that can result in neurological dysfunction, [including](#) migraine, cognitive deficit, confusion, slowed reaction times, personality changes, drowsiness and emotional changes. Some people also suffer long-term problems with memory, thinking and other symptoms, such as anxiety and mood disturbances.

After [brain injury](#) there is a cascade of events that impacts the health of neurons and affects the flow of chemical ions, such as calcium, in the brain. Mutations in genes that affect the transport of neuronal ions (atoms or molecules with a positive or negative electrical charge), termed [ion channel genes](#), can also affect how the brain works.

The strongest evidence of a connection between concussion response and ion channel gene function is from patients with a family history of a rare type of migraine (hemiplegic migraine, which causes the sufferer to experience severe migraine associated with motor impairment and muscle weakness) and [episodic ataxia](#) (which causes bouts of movement incoordination).

Specific types of these severe neurogenetic disorders are caused by mutations in the calcium channel gene [CACNA1A](#). Patients with these mutations can be highly sensitive to head impacts. Some [specific mutations](#) can see very minor head trauma lead to concussion, seizures, cerebral oedema (swelling), coma and [sometimes death](#).

Research has also shown 35% of patients with [mutations](#) in a second hemiplegic migraine ion channel gene, ATP1A2—which is linked to hemiplegic migraine, ataxia, epilepsy and other seizures and controls brain sodium and potassium levels, [report](#) concussion symptoms following mild [head trauma](#).

Focusing on all ion channel genes, our genomics lab ([Griffiths Centre for Genomics and Personalised Health](#)) recently studied 117 concussion-affected people. We found mutations in 21 ion channel genes, 14 of which could have an impact on concussion susceptibility or outcomes.

Other types of genes

Apart from a role for ion channel genes, there have been a number of additional genes linked by research to concussion.

One of the most studied is the [ApoE gene](#), which is involved in transporting cholesterol in the body and has long been recognized as a risk factor for Alzheimer's disease. Studies have indicated a variant of this gene (ApoE4) is linked with [poorer](#) and more [long-term concussion](#)

[outcomes](#). Those who carry this variant are also more likely to have significant [signs](#) of brain degeneration after concussion.

Another [genetic variation](#) in the ApoE gene that makes it less productive has been [linked](#) to a higher likelihood of concussion.

Beyond ApoE, genes that help control a variety of brain functions have been [suggested](#) as factors in concussion—including some [involved](#) in neuronal growth, dopamine receptors and, [most recently](#), brain axon (nerve fiber) development.

A predisposition for injury

Questions concerning the link between [genetic predisposition](#) to injury in sport are not new. Twenty years ago, the Australian Law Reform Commission [referred](#) to research showing "[...] a milder form of this condition [CTE or punch-drunk syndrome] could occur in players of rugby, soccer and other sports associated with repetitive blows to the head."

In 2016, the Australian Institute of Sport (AIS) released a [position statement](#) on the ethics of genetic testing and research in sport. But the [latest](#) AIS Concussion and Brain Health Position Statement does not mention the use of genetic information concerning concussion-related susceptibility.

Currently, there is available DNA diagnostic testing for the two ion channel [genes](#) already implicated in concussion, because this testing is used for the diagnosis of familial hemiplegic migraine and episodic ataxia. But genetic testing is not currently undertaken for concussion.

In Australia, it is difficult to find information on whether genetic testing occurs in elite sport. In the United Kingdom, genetic testing [does take](#)

[place](#), although it is not common. Athletes and support staff there are open to the idea of genetic information being used to improve sport performance and reduce injury risk.

What's next?

It is vital there is more careful consideration of genetic factors involved in concussion development and response. Clarification of the role of ion channel gene mutations and other gene variants, along with information from additional biomarkers and imaging, will be important in developing better concussion management and treatment approaches.

Before introducing genetic testing, regulatory and governance frameworks would also need careful consideration. Wider ethical and legal implications will need to be fully examined including health privacy laws, privacy of genetic samples, anti-discrimination laws and employment-related laws, especially in professional sport.

With the growing awareness of concussion-related injury risks highlighted by the Senate inquiry, further research in Australia could also investigate attitudes toward the use of genetic testing and predisposition to injury risk in sport.

This article is republished from [The Conversation](#) under a Creative Commons license. Read the [original article](#).

Provided by The Conversation

Citation: Genetics and concussion—why a minor knock can be devastating for some people (2023, June 20) retrieved 28 April 2024 from <https://medicalxpress.com/news/2023-06-genetics-concussionwhy-minor-devastating-people.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.