

New link between sleep and concussion identified by researchers

June 20 2023



Credit: Flinders University

A new study led by Flinders University points to the importance of sleep quality in the aftermath and recovery process in sports-related concussion, a common form of mild traumatic brain injury.

Results from a study published in *Nature and Science of Sleep* found that sleep appears to "improve" in the eight weeks after [concussion](#), including longer duration sleep, better sleep efficiency, and even longer deep sleep as the recovery progresses.

"Little is known about sleep after a concussion, despite sleep arguably being the more important process to allow the [brain](#) to function optimally," says Flinders University sleep researcher Dr. David Stevens, who treats and researches sports-related concussion.

The results of this small-scale study gives credence to previous studies in animals showing that improving sleep supports recovery from [brain injury](#), and therefore should be explored and monitored in detail for recovery and other effects of concussion such as loss of cognitive function, alertness and depression and anxiety.

"We hypothesize sleep can be a key indicator of the the brain trying to repair/recover from the concussion," the researchers note.

Previous studies have linked [deep sleep](#) leads to neural plasticity, and activation of the glymphatic system that removes amyloid- β and τ -proteins, both of which are implicated in Chronic Traumatic Encephalopathy (CTE), a progressive and fatal brain disease associated with repeated traumatic brain injuries, including concussions and repeated blows to the head.

The new study is based on evidence collected from athletes who experienced a sports-related concussion and underwent overnight sleep studies within seven days of the concussion (acute stage) and again eight weeks after the concussion (sub-acute stage).

This study indicates better sleep in participants with a concussion compared to the population norm. Sleep also appeared to further

improve in the eight weeks after concussion compared to sleep immediately after concussion.

"We speculate that the improvement in sleep was an attempt by the brain to heal itself," Dr. Stevens says.

"More research in a larger population is needed to explore this hypothesis, and investigate other neurophysiological and neurocognitive measures to examine whether changes in [sleep](#) during recovery from concussion results in changes to other aspects of the head trauma.

"I am not only advocating for improved research in a range of areas in concussion, but also for players, clubs and organizations to stop treating concussion likes it's a scratch or a 'corky,' but rather a clinical issue that has potential ramifications worse than any dislocation or fracture."

Former SANFLW player Maya Rigter says she is still recovering from three concussions, the last of which forced her to retire on 19 March.

After 57 games with Sturt Football Club and "still loving football," she decided to retire "due to the significant risk another concussion poses to my professional medical career and future health."

Currently studying physiotherapy, Rigter says the decision to stop playing "broke my heart" but ongoing symptoms were having a significant impact on her life, "particularly university with cognitive ability slow to return."

More information: David J Stevens et al, Electroencephalographic Changes in Sleep During Acute and Subacute Phases After Sports-Related Concussion, *Nature and Science of Sleep* (2023). [DOI: 10.2147/NSS.S397900](https://doi.org/10.2147/NSS.S397900)

Provided by Flinders University

Citation: New link between sleep and concussion identified by researchers (2023, June 20)
retrieved 12 May 2024 from <https://medicalxpress.com/news/2023-06-link-concussion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.