


Surgical stabilization of odontoid fractures shown to improve outcomes

June 29 2023



Two illustrative cases. Case A is that of a middle-aged woman injured in a motor vehicle collision, found on trauma workup to have A1, a nondisplaced type II

odontoid fracture that was treated with a rigid cervical orthosis and showed progressive healing at A2, 4 months and A3, 7 months after the injury. Case B is that of a middle-aged man injured in a cycling accident who had neck pain with movement and was neurologically intact on examination. CT revealed B1, a mildly displaced and angulated type II odontoid fracture treated with B2, C1-C2 posterior fixation with B3, excellent bony healing on follow-up CT. Credit: *Neurosurgery* (2023). DOI: 10.1227/neu.0000000000002557

Odontoid fractures—those occurring in the second cervical vertebra—are common in elderly patients after a low-energy fall. However, whether the initial treatment should be surgical or non-operative still isn't known. Previous studies haven't accounted for differences in injury severity, or the presence or absence of neurologic impairment, which can affect patients' results.

On this topic, an article titled "Surgery Decreases Nonunion, Myelopathy, and Mortality for Patients with Traumatic Odontoid Fractures: A Propensity Score Matched Analysis" is published in the journal *Neurosurgery*.

Michael B. Cloney, MD, MPH, of the Department of Neurological Surgery at Northwestern University in Chicago, and colleagues have published evidence that surgery should be considered as the initial approach for many patients. Compared with non-operative approaches to treatment, surgical stabilization of the fracture was associated with less myelopathy (mobility impairment due to spinal cord damage), and lower rates of fracture non-union, [30-day mortality](#), and one year mortality.

"Given the increasing incidence of odontoid fractures with the [aging population](#), we believe our findings could assist with neurosurgical decision-making for an increasingly common and complex problem," the researchers say.

Propensity score matching: A way to account for nonrandomized patient groups

Dr. Cloney and his colleagues reviewed [initial treatment](#) data on 296 patients who were cared for at Northwestern Memorial Hospital between January 1, 2010, and December 31, 2020, because of an odontoid fracture. Their average age was 73. During the hospitalization, 22% had surgery and 78% had non-operative treatment (5% were immobilized in a halo-vest and 73% received a cervical collar).

Since the patients weren't randomized to these treatments, the research team used a type of analysis called propensity score adjustment. They calculated "propensity scores" for each individual—the probability that the patient would have been assigned to receive one of the two treatment approaches based on certain characteristics.

For example, to study the effect of surgery on [mortality rates](#), patients were matched on age, sex, Injury Severity Score, Nurick score (a measure of myelopathy), their number of chronic diseases and chronic conditions such as smoking, and whether they had to be admitted to the intensive care unit.

Surgical stabilization leads to better results

Follow up with patients lasted an average of 45 weeks. On the [propensity score](#)-matched analyses, the group that underwent surgery showed significantly better outcomes than the non-operative group:

- Lower rate of fracture non-union—39.7% vs. 57.3%; treatment effect, 15% less risk of nonunion
- Lower 30-day mortality rate—1.7% vs. 13.8%; treatment effect, 10% less risk of death

- Lower one year mortality rate—7.0% vs. 23.7%; treatment effect, 10% less risk of death

Other analyses showed patients in the surgery group were 52% less likely than those in the non-operative group to have poor Nurick scores at the 26-week postoperative follow-up visit and were 41% less likely to die during the overall follow-up period. Both differences were statistically significant.

"The mortality benefit calculated in the existing literature typically represents an unadjusted mortality rate between two potentially different populations, which leaves it liable to confounding," the authors note.

"Our study represents a relatively large institutional series that suggests a benefit from surgical stabilization in this population while controlling for confounding factors more thoroughly than existing literature."

More information: Michael Cloney et al, Surgery Decreases Nonunion, Myelopathy, and Mortality for Patients With Traumatic Odontoid Fractures: A Propensity Score Matched Analysis, *Neurosurgery* (2023). [DOI: 10.1227/neu.0000000000002557](https://doi.org/10.1227/neu.0000000000002557)

Provided by Wolters Kluwer Health

Citation: Surgical stabilization of odontoid fractures shown to improve outcomes (2023, June 29) retrieved 27 April 2024 from

<https://medicalxpress.com/news/2023-06-surgical-stabilization-odontoid-fractures-shown.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.