

Why women with multiple sclerosis get better when pregnant

June 13 2023



Credit: CC0 Public Domain

Women suffering from the autoimmune disease multiple sclerosis temporarily get much better when pregnant. Researchers have now identified the beneficial changes naturally occurring in the immune system during pregnancy. The findings, published in *Journal of Neuroinflammation*, may pave the way to new treatments.

Pregnancy is a very special condition from an immunological point of view. The [immune system](#) serves to defend us against foreign substances. However, although half of the genetic material of the fetus comes from the father, it is not rejected by the mother's immune system. One reason why this balancing act is almost always successful is that during pregnancy the mother's immune system is adapted to become more tolerant.

In multiple sclerosis, MS, nerve function is hampered due to the immune system attacking the fat that serves as an insulating sheath around the [nerve fibers](#). The nerves become inflamed, which could lead to nerve damage. Although new and more effective treatment options are available, most MS patients deteriorate over time.

Researchers believe that the temporary dampening of the immune response could explain why women with MS actually get better when pregnant. Periods of symptoms, i.e. relapses, decrease by 70% during the third trimester. Also some other autoimmune diseases, such as rheumatoid arthritis, temporarily ameliorate during pregnancy. But the reason for this has not been clear. This is why the researchers wanted to investigate mechanisms that could be of particular importance in the reduction of symptoms during pregnancy, as a step to finding future treatment strategies that give the same effect in MS and possibly also other similar diseases.

The researchers were particularly interested in T cells, which play an important role in the immune system. Moreover, T cells play a key role in driving MS and are important during pregnancy. The study compared 11 women with MS to seven healthy women who had blood samples taken before, during and after pregnancy.

To understand what happens in the immune cells, the researchers identified the genes used in the T cells at various points in time during

pregnancy. They also studied changes regulating how genes are switched on and off, i.e. epigenetic changes. In their study, the researchers looked more specifically at one such regulation mechanism called DNA methylation.

"What was possibly most striking is that we couldn't find any real differences between the groups during pregnancy, as it seems that the immune system of a pregnant woman with MS looks roughly like that of a healthy pregnant woman," says Sandra Hellberg, assistant professor at the Department of Biomedical and Clinical Sciences at Linköping University and one of the researchers behind the study.

The researchers found networks of interacting genes that are affected during pregnancy. Their study shows that these genes are to a large extent linked to the disease and to important processes in the immune system.

"We can see that the changes in the T cells mirror the amelioration in relapse frequency. The biggest changes happen in the last third of pregnancy, and this is where women with MS improve the most. These changes are then reversed after pregnancy at the point in time when there is a temporary increase in disease activity. It is important to stress that disease activity thereafter goes back to what it was prior to the pregnancy," says Sandra Hellberg.

The network of genes affected during pregnancy also included genes regulated by [pregnancy](#) hormones, mainly progesterone. The researchers are now testing various hormones in the lab in an attempt to mimic the effects observed in the study, to see if these can be part of a possible future treatment strategy.

This research is the result of long-standing collaboration between researchers in medicine and bioinformatics. A key part of the project

has been understanding the large amount of data by analyzing it using what is known as [network analysis](#), developed over many years by, among others, a research group led by Mika Gustafsson at Linköping University. Network analysis is a tool for finding genes that interact extensively with the genes the researchers are interested in. It often turns out that other [genes](#) in the network are regulated in an abnormal manner and indirectly affect key processes in a disease.

"Such insights can be used to find alternative medication and find new biomarkers to be able to differentiate between subgroups of a disease. We have used this strategy successfully for analysis in research into for instance allergy and multiple sclerosis," says Mika Gustafsson, professor of Bioinformatics, who is now making the analysis available to other researchers through a newly founded company.

The study is a sub-study of the GraMS (Pregnancy and MS) study and was carried out in collaboration between Linköping University, the Karolinska University Hospital in Solna, Linköping University Hospital, Länssjukhuset Ryhov in Jönköping and Region Kalmar.

More information: Alberto Zenere et al, Prominent epigenetic and transcriptomic changes in CD4+ and CD8+ T cells during and after pregnancy in women with multiple sclerosis and controls, *Journal of Neuroinflammation* (2023). [DOI: 10.1186/s12974-023-02781-2](https://doi.org/10.1186/s12974-023-02781-2)

Provided by Linköping University

Citation: Why women with multiple sclerosis get better when pregnant (2023, June 13) retrieved 29 April 2024 from

<https://medicalxpress.com/news/2023-06-women-multiple-sclerosis-pregnant.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.