



Brain movement measured for clues to prevent, reduce injury

July 27 2023, by Beth Miller



Tagged MRI methods. (a) and (b) The participant uses a neck rotation device to obtain repeatable, mild, impulsive motion. (a) The participant in rest position before rotation. (b) The participant activates a latch that allows head rotation toward the left shoulder until motion is arrested by a compliant stop. Images adapted from Gomez et al. [38]. (c) and (d) Schematic depiction of head motion. (e) Representative anatomical (T1-weighted) axial MR image of the participant's brain. (f) Representative tagged MR image of the brain approximately 50 ms

after impact (adapted from Escarcega et al. [32]). Tag lines are initially straight and move with the tissue, revealing deformation. Scale bar is 5 cm. Credit: *Journal of Biomechanical Engineering* (2023). DOI: 10.1115/1.4062809

When the human head experiences any kind of movement—from nodding yes or no to heading a soccer ball or being jolted in a car crash—the brain moves inside the skull, leading to deformation of the tissue. Such deformations are key to understanding traumatic brain injury but are challenging to study since the brain is hidden inside the skull.

Philip V. Bayly, the Lee Hunter Distinguished Professor and chair of the Department of Mechanical Engineering & Materials Science at the McKelvey School of Engineering at Washington University in St. Louis, and Jordan D. Escarcega, a [mechanical engineering](#) doctoral student in Bayly's lab, led a multi-institutional team to compare how the human brain deforms in response to movement using two types of magnetic resonance imaging (MRI). Their work is published in the *Journal of Biomechanical Engineering*.

The researchers measured the impulse response of the brain—its natural reaction to impact—and its harmonic response, or response to skull vibration at a particular frequency, similar to the experience of driving on rumble strips on the side of the road.

In the first case, the human volunteers' heads were resting in a cradle while undergoing tagged MRI imaging. The volunteers were asked to gently rotate their head side to side—as if shaking their head "no"—until the cradle stopped the movement, which created the impulse response. To see the impulse response of the brain in tagged MRI, a grid of lines is superimposed over the MR image of the moving brain to see the

deformation of the brain when the volunteer's head was stopped by the cradle.

In the second experiment, light sound pressure generated by a speaker was applied to the volunteers' heads to generate a harmonic response. Brain motion was measured using MR elastography, or MRE, a noninvasive technique that combines MRI images with low-frequency vibrations to create a visual map that shows information on body tissues.

"In the harmonic response, you're getting a single sustained wave pattern, whereas in an impulse response, you're getting a complex behavior that consists of the responses to multiple frequencies added together," said Escarcega, who first came to Washington University as part of the Washington University Summer Engineering Fellowship Program (WUSEF), then returned to earn a doctorate.

"Using the tagged MRI, we are able to break down the impulse response into the most dominant components and compare those to the harmonic waveforms that we excite in MRE."

"Every [human brain](#) is different—it might as well be like your fingerprint," Escarcega said. "But what we found was the physical patterns of deformation produced by MR elastography are extremely similar to the patterns you see from when people bump their heads."

Tagged MRI is more difficult to perform than MRE because it requires participants to move their heads many times. Bayly and Escarcega said that MRE, which is simpler and more widely available, can be used to show the brain's response to head impact. This result has important practical implications, they said.

"Using MRE, we can quantitatively measure patterns of brain deformation that are expected in response to head impact, but without

the longer experiments and added difficulty of tagged MRI," Bayly said. "These measurements will be useful to evaluate and improve computer models of brain biomechanics, which can then inform the design of protective equipment and reduce the societal burden of [traumatic brain injury](#)."

More information: Jordan D. Escarcega et al, Comparison of Deformation Patterns Excited in the Human Brain In Vivo by Harmonic and Impulsive Skull Motion, *Journal of Biomechanical Engineering* (2023). [DOI: 10.1115/1.4062809](https://doi.org/10.1115/1.4062809)

Provided by Washington University in St. Louis

Citation: Brain movement measured for clues to prevent, reduce injury (2023, July 27) retrieved 9 May 2024 from <https://medicalxpress.com/news/2023-07-brain-movement-clues-injury.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--